

4 - OCT 3 1
Copy / 1962



Radiological Health Data

VOLUME III, NUMBER 10

OCTOBER 1962

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

Public Health Service

In August 1959, the President directed the Secretary of Health, Education, and Welfare to intensify Departmental activities in the field of radiological health. The Department was assigned responsibility within the Executive Branch for the collation, analysis and interpretation of data on environmental radiation levels. The Department delegated this responsibility to the Division of Radiological Health, Public Health Service.

Radiological Health Data is published by the Public Health Service on a monthly basis. Data are provided to the Division of Radiological Health by other Federal agencies, State health departments, and foreign governments. Except where material is directly quoted or otherwise credited, summaries and abstracts are prepared by the Radiological Health Data and Reports Staff, Division of Radiological Health. The reports are reviewed by a Board of Editorial Advisors with representatives from the following Federal agencies:

Department of Health, Education, and
Welfare
Atomic Energy Commission
Department of Defense
Department of Agriculture
Department of Commerce

For further information on any subject reported in this issue, readers are referred to the contributors indicated in the headings of the articles.

For personal and extra subscriptions to *Radiological Health Data*, use the handy order form on inside back cover.

—
T

Ad

SE

Fi

Na

St

St


De

St

Ra

Ra

—
O





RADIOLOGICAL HEALTH DATA

VOLUME III, NUMBER 10
OCTOBER 1962

TABLE OF CONTENTS

| | Page | | Page |
|--------------------------------------------------------------------------------------------------------------------------------------|------|-----------------------------------------------------------------------------------------------|-------------------|
| Advance Report | ii | SECTION III.—MILK—Continued | |
| SECTION I.—AIR AND PRECIPITATION | | Strontium-90 in Canadian Dried Milk Products (First Quarter 1962) | |
| Fission Product Beta Activity in Airborne Particu- lates and Precipitation | 379 | Iodine-131 and Strontium-90 in Milk From the United Kingdom (January 1961–April 1962) | 404 |
| Radiation Surveillance Network (July 1962), PHS | | SECTION IV.—WATER | |
| The 80th Meridian (West) Sampling Program (June 1962), NRL | | Radioactivity in Raw Surface Waters (April 1962), PHS | |
| National Air Sampling Network (Second Quarter 1962), PHS | 385 | Radionuclide Analyses of Coast Guard Water Sup- plies (October 1961–April 1962), PHS | 410 |
| Canadian Radioactive Fallout Study Program (First Quarter 1962) | 388 | Radioactivity in Drinking Water (1961), PHS | 411 |
| Natural and Fission Product Radioactivity in Sur- face Air Particulates at Cincinnati, Ohio (June 13–July 13, 1962), PHS | 389 | SECTION V.—OTHER DATA | |
| SECTION II.—FOOD | | Whole Body Counting | |
| Strontium-90 Concentrations in U.S. Wheat and Milling Products (1960), AEC | 392 | Cesium-137 in Man (March–May 1962), Land- stuhl, Germany | 413 |
| Deposition of Strontium-90 in a 1960 Kansas Wheat Sample, AEC | 395 | External Gamma Activity (July 1962), PHS | 414 |
| SECTION III.—MILK | | Environmental Levels of Radioactivity at Atomic Energy Commission Installations | 415 |
| Radionuclides in Pasteurized Milk (June 1962), PHS | 397 | Hanford Atomic Products Operation (Third and Fourth Quarters 1961) | 416 |
| Radionuclides in Raw Milk From the Vicinity of the Nevada Test Site (July 1961–June 1962), AEC | 402 | Lawrence Radiation Laboratory (Third and Fourth Quarters 1961) | 418 |
| | | Los Alamos Scientific Laboratory (1960–1961) | 419 |
| | | Nevada Test Site Detonation Surveillance | 420 |
| | | Reported Nuclear Detonations (September 1962) .. | 423 |
| | | Announcement of Current Training Activities PHS Units and Equivalents | Inside Back Cover |

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service • Division of Radiological Health

ADVANCE REPORT

Reports received by the Public Health Service Radiation Surveillance Center for August and September 1962 have indicated fresh fallout from both U. S. and U.S.S.R. testing. The Service's Pasteurized Milk Network reports show daily levels of radioactive iodine-131 from July Nevada tests declined in Salt Lake City milk from 730 to 100 $\mu\mu\text{C}$ per liter in August. The August Salt Lake City monthly average was 330 $\mu\mu\text{C}/\text{l}$. The only other station with an iodine-131 August monthly average over 40 $\mu\mu\text{C}/\text{l}$ was Laramie, Wyoming with 160 $\mu\mu\text{C}/\text{l}$. The September monthly iodine-131 averages for both Salt Lake City, Utah and Laramie, Wyoming were 40 $\mu\mu\text{C}/\text{l}$.

Apparent effects of the U.S.S.R. nuclear tests appeared in September when the Pasteurized Milk Network monthly tabulation of 62 stations showed an average daily iodine-131 concentration of 60 $\mu\mu\text{C}/\text{l}$ compared to the August average of 20 $\mu\mu\text{C}/\text{l}$. During September the monthly averages of 40 stations were above 20 $\mu\mu\text{C}/\text{l}$ compared to 6 stations in August. Of these 40, the following 9 stations, all in the northern States, had September monthly averages of 100 $\mu\mu\text{C}/\text{l}$ or more:

| | $\mu\mu\text{C}/\text{l}$ | | $\mu\mu\text{C}/\text{l}$ |
|-------------------------|---------------------------|--------------------------------|---------------------------|
| Palmer, Alaska | 730 | Buffalo, New York | 100 |
| Chicago, Illinois | 100 | Minot, North Dakota | 170 |
| Des Moines, Iowa | 120 | Rapid City, South Dakota | 130 |
| Detroit, Michigan | 120 | Milwaukee, Wisconsin | 120 |

SECTION I.—AIR AND PRECIPITATION

Fission Product Beta Activity in Airborne Particulates and Precipitation

Measurements of gross beta activity of airborne particulates and precipitation are among the earliest and most sensitive indicators of increases of fission product activity in the environment. However, a direct evaluation of biological effects is not possible from these data alone.

Of the several networks or sampling programs making such measurements, the Radiation Surveillance Network, the Naval Research Laboratory, the National Air Sampling Network, and the Canadian Radioactive Fallout Study Program are represented in the following reports.

RADIATION SURVEILLANCE NETWORK July 1962

Division of Radiological Health, Public Health Service

The Public Health Service Radiation Surveillance Network (RSN) was established in 1956 in cooperation with the Atomic Energy Commission primarily to provide a means of promptly determining increasing levels of radioactivity in air and precipitation due to fallout from nuclear weapons tests. Prior to September 1961, the Network consisted of 45 stations. Following the September 1961 resumption of nuclear weapons testing by the U.S.S.R., the Network has been expanded over

a period of several months to 72 stations, whose locations are shown in figure 1.

Air

Daily 24-hour air samples are collected by a high volume air sampler with a 4-inch diameter carbon-loaded cellulose dust filter. Field measurements with a portable survey meter enable the station operator to estimate the amount of beta activity in airborne particulates at the station five hours after collection by comparison with a known $\text{Sr}^{90}\text{--Y}^{90}$ source. This 5-hour delay eliminates interference from naturally-occurring radon daughters. Each operator then reports his field estimate by telephone to the Radiation Surveillance Center, Division of Radiological Health, Washington, D.C., to provide a daily national report.

The filters are then forwarded to the Radiation Surveillance Network Laboratory in Rockville, Maryland, for a more refined measurement using a thin-window gas-flow proportional counter. Each filter is counted at least 3 days after the end of the sampling period and recounted 7 days later. The initial 3-day aging of the sample eliminates interference from naturally-occurring radon and thoron daughters. The two counts, separated by a 7-day interval, make possible the estimation of the age of fission products and extrapolation of the




FIGURE 1.—RADIATION SURVEILLANCE NETWORK SAMPLING STATIONS AND AVERAGE FISSION PRODUCT BETA CONCENTRATIONS IN AIR ($\mu\text{pc}/\text{m}^3$) JULY 1962

activity to the time of collection. The extrapolation is performed by using the Way-Wigner formula ($AT^{1.2} = \text{constant}$).*

Although gross beta concentration is usually presented without reference to age, it is apparent that fission product activity is more adequately described when the age parameter is also given. An example of the interpretative usefulness of age of fission product data is presented in "Mid-May Iodine-131 in Fallout in the Midwest" in *RHD*, September 1962, Section V. Because of the difficulty in determining a meaningful monthly average of age of fission products, such data have not yet been presented with the RSN data in *RHD*.

The average fission-product beta concentrations in surface air during July 1962 as determined by laboratory analysis are tabulated in table 1 and presented by means of concentration contours in figure 1. Experience has shown that field estimates are generally comparable to laboratory analyses, except at low levels, where

* In this expression, A is the activity at time T after fission product formation. Units are arbitrary.

the former are usually higher because of natural thoron daughters.

Precipitation

Continuous sampling for total precipitation is conducted at most stations on a daily basis using funnels having collection areas of 0.4 m^2 . A 500 ml aliquot of the collected precipitation is evaporated to dryness, and the residue is forwarded to the laboratory to be counted by the same method used for analyzing the air samples. If the collected sample is between 200 and 500 ml, the entire sample is evaporated; if less than 200 ml (equivalent to 0.5 mm rainfall), the volume of precipitation is reported, but no analysis is made.

The June 1962 averages of gross beta activity in precipitation, expressed in micromicrocuries per liter ($\mu\mu\text{c}/\text{liter}$) and millimicrocuries per square meter ($\text{m}\mu\text{c}/\text{m}^2$) are presented in table 2. Placement of a "less than" sign (<) with an average concentration or total deposition value indicates that the sum of the "less than" daily deposition values is 10 percent or more of the

TABLE 1.—GROSS BETA ACTIVITY OF PARTICULATES
IN AIR, RSN, JULY 1962[Concentrations in $\mu\text{mc}/\text{m}^3$]

| Station location | | Number samples | Maximum ($\mu\text{mc}/\text{m}^3$) | Minimum ($\mu\text{mc}/\text{m}^3$) | Average* |
|------------------|---------|-------------------|------------------------------------------|------------------------------------------|----------|
| City | State | | | | |
| Adak | Alaska | 24 | 1.6 | <0.10 | 0.40 |
| Anchorage | Alaska | 31 | 5.3 | <0.10 | 1.1 |
| Attu | Alaska | 30 | 3.2 | <0.10 | 0.50 |
| Fairbanks | Alaska | 16 | 2.7 | <0.10 | 1.6 |
| Juneau | Alaska | 19 | 3.4 | 0.47 | 1.0 |
| Kodiak | Alaska | 31 | 2.9 | <0.10 | 0.71 |
| Nome | Alaska | 28 | 4.7 | <0.10 | 0.93 |
| Point Barrow | Alaska | 28 | 1.8 | <0.10 | 0.43 |
| St. Paul Island | Alaska | 16 | 1.1 | <0.10 | 0.31 |
| Phoenix | Ariz. | 30 | 14 | 1.2 | 3.1 |
| Little Rock | Ark. | 31 | 12 | 1.1 | 3.3 |
| Berkeley | Calif. | 31 | 2.5 | 0.17 | 1.2 |
| Los Angeles | Calif. | 23 | 3.1 | 0.84 | 2.0 |
| Denver | Colo. | 31 | 93 | 1.3 | 7.3 |
| Hartford | Conn. | 31 | 2.5 | 0.61 | 2.5 |
| Dover | Del. | 22 | 7.1 | 1.5 | 4.1 |
| Washington | D. C. | 31 | 6.9 | 0.81 | 2.9 |
| Jacksonville | Fla. | 31 | 5.5 | 0.72 | 2.4 |
| Miami | Fla. | 31 | 3.4 | 0.90 | 1.6 |
| Atlanta | Ga. | 31 | 4.8 | 0.47 | 2.5 |
| Agana | Guam | 13 | 19 | <0.10 | 2.9 |
| Honolulu | Hawaii | 29 | 2.5 | 0.18 | 1.1 |
| Boise | Idaho | 28 | 25 | 2.1 | 4.2 |
| Springfield | Ill. | 31 | 6.5 | 1.1 | 2.5 |
| Indianapolis | Ind. | 30 | 4.5 | 0.83 | 2.4 |
| Iowa City | Iowa | 31 | 3.9 | 0.69 | 1.9 |
| Topeka | Kans. | 31 | 5.9 | 0.94 | 2.3 |
| Frankfort | Ky. | 29 | 4.9 | 0.93 | 2.1 |
| New Orleans | La. | 31 | 5.7 | 0.82 | 2.5 |
| Augusta | Maine | 31 | 5.1 | 0.12 | 2.3 |
| Presque Isle | Maine | 25 | 3.7 | 0.15 | 1.4 |
| Baltimore | Md. | 31 | 6.6 | 1.2 | 3.2 |
| Rockville | Md. | 22 | 6.5 | 0.90 | 3.3 |
| Lawrence | Mass. | 31 | 4.4 | 0.16 | 2.0 |
| Winchester | Mass. | 28 | 4.5 | 0.17 | 2.3 |
| Lansing | Mich. | 31 | 6.0 | 1.7 | 3.5 |
| Minneapolis | Minn. | 31 | 5.6 | 0.80 | 2.3 |
| Jackson | Miss. | 27 | 12 | 0.43 | 3.7 |
| Pascagoula | Miss. | 21 | 8.9 | 1.0 | 3.8 |
| Jefferson City | Mo. | 31 | 8.1 | 1.1 | 2.8 |
| Helena | Mont. | 31 | 7.4 | 0.79 | 3.7 |
| Lincoln | Nebr. | 21 | 5.5 | 0.96 | 2.7 |
| Las Vegas | Nev. | 30 | 17 | 0.59 | 6.6 |
| Concord | N. H. | 21 | 5.9 | 0.29 | 2.9 |
| Trenton | N. J. | 31 | 4.6 | 1.1 | 2.7 |
| Santa Fe | N. Mex. | 31 | 8.7 | 1.2 | 2.9 |
| Albany | N. Y. | 30 | 5.1 | 0.34 | 2.8 |
| Buffalo | N. Y. | 30 | 6.1 | 1.8 | 3.3 |
| New York | N. Y. | 17 | 4.6 | 0.92 | 2.4 |
| Gastonia | N. C. | 31 | 7.9 | 0.84 | 3.5 |
| Bismark | N. Dak. | 31 | 8.6 | 0.71 | 3.2 |
| Columbus | Ohio | 30 | 6.5 | 0.99 | 2.8 |
| Painesville | Ohio | 30 | 5.5 | 1.2 | 3.0 |
| Oklahoma City | Okla. | 31 | 5.1 | 0.76 | 1.9 |
| Ponca City | Okla. | 29 | 3.2 | 0.12 | 1.1 |
| Portland | Ore. | 31 | 3.0 | 0.64 | 1.6 |
| Harrisburg | Pa. | 24 | 6.3 | 0.28 | 3.4 |
| San Juan | P. R. | 15 | 2.5 | 0.84 | 1.5 |
| Providence | R. I. | 31 | 5.9 | 0.18 | 2.5 |
| Columbia | S. C. | 27 | 4.9 | 0.40 | 2.6 |
| Pierre | S. Dak. | 30 | 40 | 0.85 | 4.0 |
| Nashville | Tenn. | 30 | 9.1 | 1.1 | 3.4 |
| Austin | Tex. | 29 | 4.2 | 0.68 | 1.6 |
| El Paso | Tex. | 30 | 3.9 | 0.66 | 1.4 |
| Salt Lake City | Utah | 31 | 595 | 1.7 | 51 |
| Barre | Vt. | 31 | 5.8 | 0.33 | 3.0 |
| Richmond | Va. | 31 | 4.0 | 0.78 | 2.4 |
| Seattle | Wash. | 30 | 2.2 | 0.47 | 0.95 |
| Charleston | W. Va. | 29 | 4.6 | 1.2 | 2.6 |
| Madison | Wis. | 31 | 5.1 | 1.2 | 3.0 |
| Cheyenne | Wyo. | 29 | 378 | 1.3 | 21 |

Network average..... 3.4

* Weighted average obtained by summing the products of individual sampling times and the corresponding activities, and dividing by the summation of the individual sampling times.

TABLE 2.—GROSS BETA ACTIVITY IN
PRECIPITATION, RSN, JULY 1962

| Station location | | Average concentration ($\mu\text{mc}/\text{liter}$) | Total deposition ($\text{m}\mu\text{c}/\text{m}^2$) |
|------------------|---------|-------------------------------------------------------------|-------------------------------------------------------------|
| City | State | | |
| Adak | Alaska | * | * |
| Anchorage | Alaska | 610 | 6 |
| Attu | Alaska | * | * |
| Fairbanks | Alaska | * | * |
| Juneau | Alaska | 650 | 55 |
| Kodiak | Alaska | * | * |
| Nome | Alaska | * | * |
| Point Barrow | Alaska | * | * |
| St. Paul Island | Alaska | * | * |
| Phoenix | Ariz. | * | * |
| Little Rock | Ark. | 410 | 39 |
| Berkeley | Calif. | * | * |
| Los Angeles | Calif. | * | * |
| Denver | Colo. | * | * |
| Hartford | Conn. | 2,000 | 75 |
| Dover | Del. | * | * |
| Washington | D. C. | 2,500 | 85 |
| Jacksonville | Fla. | 510 | 69 |
| Miami | Fla. | 1,700 | 180 |
| Atlanta | Ga. | † <360 | † <24 |
| Agana | Guam | * | * |
| Honolulu | Hawaii | * | * |
| Boise | Idaho | * | * |
| Springfield | Ill. | 2,200 | 120 |
| Indianapolis | Ind. | 790 | 160 |
| Iowa City | Iowa | 500 | 130 |
| Topeka | Kans. | 1,200 | 83 |
| Frankfort | Ky. | 700 | 48 |
| New Orleans | La. | 990 | 90 |
| Augusta | Maine | 640 | 43 |
| Presque Isle | Maine | 560 | 91 |
| Baltimore | Md. | 930 | 55 |
| Rockville | Md. | * | * |
| Lawrence | Mass. | 680 | 69 |
| Winchester | Mass. | 710 | 31 |
| Lansing | Mich. | 620 | 25 |
| Minneapolis | Minn. | 2,100 | 280 |
| Jackson | Miss. | † <350 | † <6.5 |
| Pascagoula | Miss. | 550 | 110 |
| Jefferson City | Mo. | 2,600 | 53 |
| Helena | Mont. | 7,400 | 300 |
| Lincoln | Nebr. | 2,200 | 240 |
| Las Vegas | Nev. | * | * |
| Concord | N. H. | * | * |
| Trenton | N. J. | 1,400 | 3.3 |
| Santa Fe | N. Mex. | 990 | 52 |
| Albany | N. Y. | 660 | 15 |
| Buffalo | N. Y. | 820 | 16 |
| New York | N. Y. | * | * |
| Gastonia | N. C. | 1,000 | 82 |
| Bismarck | N. Dak. | 1,500 | 110 |
| Columbus | Ohio | 850 | 89 |
| Painesville | Ohio | 960 | 100 |
| Oklahoma City | Okla. | 470 | 9 |
| Ponca City | Okla. | 600 | 64 |
| Portland | Ore. | * | * |
| Harrisburg | Pa. | 1,400 | 18 |
| San Juan | P. R. | 350 | 17 |
| Providence | R. I. | 650 | 23 |
| Columbia | S. C. | 610 | 81 |
| Pierre | S. Dak. | 2,100 | 94 |
| Nashville | Tenn. | 810 | 29 |
| Austin | Tex. | * | * |
| El Paso | Tex. | 980 | 57 |
| Salt Lake City | Utah | 6,700 | 140 |
| Barre | Vt. | 1,000 | 130 |
| Richmond | Va. | 460 | 59 |
| Seattle | Wash. | 630 | 5 |
| Charleston | W. Va. | 890 | 110 |
| Madison | Wis. | 1,200 | 62 |
| Cheyenne | Wyo. | 2,600 | 170 |

* No evaporated sample received.

† Placement of a "less than" sign (<) with an average concentration or total deposition value indicates that the sum of the "less than" daily values is 10 percent or more of the total, so that the true total or average is considered significantly less than the value shown.




FIGURE 2.—MONTHLY AND YEARLY PROFILES OF BETA ACTIVITY IN AIR, RADIATION SURVEILLANCE NETWORK, 1956-JULY 1962

total deposition so that the true total or average is considered significantly less than the value shown.

Profiles

The profiles of the monthly average fission product beta activity in airborne particulates for each RSN station covering the period of time from the formation of the network in 1956 to the end of 1960 were published in *RHD*, July 1961. The profiles of 7 stations, updated through July 1962, are shown in figure 2.

THE 80TH MERIDIAN (WEST) SAMPLING PROGRAM June 1962

U.S. Naval Research Laboratory

Radioactivity measurements of surface air samples collected at various sites near the 80th Meridian (West) have been made since 1956. Sampling locations are shown in figure 3. This program is operated by the U.S. Naval Research Laboratory (NRL) with the cooperation of interested agencies of the United States, Canada, Ecuador, Peru, Bolivia, and Chile, which collect the samples and forward them to NRL for analysis. Partial financial support of




FIGURE 3.—ATMOSPHERIC RADIOACTIVITY SAMPLING STATIONS NEAR THE 80TH MERIDIAN (WEST)

this program is provided by the Division of Biology and Medicine, U.S. Atomic Energy Commission.




FIGURE 4.—PROFILE OF BETA ACTIVITY, AVERAGE MEASUREMENT OF SURFACE AIR AT STATIONS NEAR THE 80TH MERIDIAN (WEST), JUNE 1962

TABLE 3.—FISSION PRODUCT GROSS BETA ACTIVITY IN SURFACE AIR, NRL,^a JUNE 1962[Average concentrations in $\mu\text{mc}/\text{m}^3$]

| Date | Punta Arenas, Chile | Puerto Montt, Chile | Santiago, Chile | Antofagasta, Chile | Chacaltaya, Bolivia | Lima, Peru | Guayaquil, Ecuador | Miraflores, Panama Canal Zone | San Juan, P. R. | Mauna Loa, Hawaii ^c | Miami, Florida | Washington, D. C. | Moosonee, Ontario, Canada | Thule, Greenland |
|------------------|---------------------|---------------------|-----------------|--------------------|---------------------|------------|--------------------|-------------------------------|-----------------|--------------------------------|----------------|-------------------|---------------------------|------------------|
| 1 | 0.018 | 0.021 | 0.159 | 1.68 | 4.22 | 2.90 | b— | 1.00 | 0.805 | 3.48 | 2.65 | 4.76 | 3.62 | 5.31 |
| 2 | | | | | | | | | | | | | | |
| 3 | | | | | | | | | | | | | | |
| 4 | | | | | | | | | | | | | | |
| 5 | 0.018 | 0.056 | 0.377 | 1.84 | 2.36 | 1.52 | — | 0.210 | 3.16 | 3.99 | 2.34 | 7.91 | 7.50 | 4.95 |
| 6 | | | | | | | | | | | | | | |
| 7 | | | | | | | | | | | | | | |
| 8 | | | | | | | | | | | | | | |
| 9 | 0.041 | 0.063 | 0.964 | 2.63 | 1.39 | 2.85 | — | 0.209 | 2.42 | 3.39 | 1.37 | 5.40 | 8.00 | 2.82 |
| 10 | | | | | | | | | | | | | | |
| 11 | | | | | | | | | | | | | | |
| 12 | | | | | | | | | | | | | | |
| 13 | 0.052 | 0.050 | — | 1.02 | 1.46 | 2.04 | — | 0.193 | 2.54 | 3.16 | 1.81 | 7.78 | 4.95 | 1.79 |
| 14 | | | | | | | | | | | | | | |
| 15 | | | | | | | | | | | | | | |
| 16 | | | | | | | | | | | | | | |
| 17 | 0.057 | 0.078 | — | — | 1.33 | 1.76 | — | 0.193 | 2.21 | 4.30 | 3.06 | 8.81 | 5.21 | 1.58 |
| 18 | | | | | | | | | | | | | | |
| 19 | | | | | | | | | | | | | | |
| 20 | | | | | | | | | | | | | | |
| 21 | 0.057 | 0.078 | — | — | 1.33 | 1.76 | — | 0.193 | 2.21 | 4.30 | 3.06 | 8.81 | 5.21 | 1.58 |
| 22 | | | | | | | | | | | | | | |
| 23 | | | | | | | | | | | | | | |
| 24 | | | | | | | | | | | | | | |
| 25 | 0.057 | 0.078 | — | — | 1.33 | 1.76 | — | 0.193 | 2.21 | 4.30 | 3.06 | 8.81 | 5.21 | 1.58 |
| 26 | | | | | | | | | | | | | | |
| 27 | | | | | | | | | | | | | | |
| 28 | | | | | | | | | | | | | | |
| 29 | 0.057 | 0.078 | — | — | 1.33 | 1.76 | — | 0.193 | 2.21 | 4.30 | 3.06 | 8.81 | 5.21 | 1.58 |
| 30 | | | | | | | | | | | | | | |
| Weighted average | 0.038 | 0.055 | 0.558 | 1.80 | 2.00 | 2.18 | — | 0.309 | 2.37 | 3.64 | 2.15 | 6.93 | 6.11 | 3.32 |

^a The average concentration determined from a given sample is placed at the center of a rectangle which indicates the length and dates of the sampling period. Station averages for the month were determined by weighting the sample average according to the number of days in the sampling period of the portion of the sampling period occurring in June 1962.

^b Dash indicates samples were not received.

^c Mauna Loa data has been included for comparison with Chacaltaya, Bolivia. Both are high elevation stations (3400 and 5200 meters) and about equally distant north and south of the equator.

The sampling procedure involves drawing air continuously for a 7-day period at a rate of approximately 1200 cubic meters per day through an 8-inch diameter, high efficiency filter, using a positive displacement blower. After the 7-day period, the filter is removed and forwarded to NRL for assay of gross beta activity. A minimum of 2 weeks after collection is allowed for decay of short-lived radionuclides. Data are not extrapolated to time of collection.

The data in table 3 and the June profile in figure 4 show a conspicuous low concentration at Miraflores compared to the adjacent stations. This was perhaps the result of two factors: (a) fresh radioactivity from U.S. Pacific tests went much further south while the Northern Hemisphere stations remained high because of the background of 1961 U.S.S.R. debris (plus perhaps some fallout from U.S. Pacific tests); and (b) the rainy season at Panama depressed the activity concentrations in the ground-level air.

NATIONAL AIR SAMPLING NETWORK Second Quarter 1962

Division of Air Pollution, Public Health Service

The Public Health Service developed its National Air Sampling Network in 1953 to secure basic data on the nature and extent of air pollution throughout the United States and to detect trends in levels of pollution with respect to time, location, population density, climate, and other factors associated with air quality.

The current basic network consists of 110 sampling stations operating every year in 73 large cities and 37 nonurban areas. In addition to these every-year stations, 130 cities have stations which operate every other year. Thus, there are 240 sampling stations in all, of which about 175 are active in any given year.

The network stations are manned by co-operating Federal, State, and local agencies.

Twenty-four hour samples of suspended particulate matter representing approximately 2000 cubic meters of air are collected on glass fiber filters on a bi-weekly random sampling schedule. The analyses of these samples include the measurement of total quantity of suspended particulate matter, the organic matter soluble in benzene, and gross beta radioactivity. Selected samples are analyzed also for nitrates, sulfates, and a number of metals.

Gross Beta Activity in Air

Gross beta activity data by states, for the years 1953 through 1958, were submitted by the Division of Radiological Health, Public Health Service, in testimony before the Joint Committee on Atomic Energy Hearing on Fallout from Nuclear Weapons Tests, Volume I, May 1959, pages 173-185. Subsequent data have been published quarterly in *Radiological Health Data* beginning with the October 1960 issue. Second quarter 1962 data are presented in table 4.




FIGURE 5.—NATIONAL AIR SAMPLING NETWORK STATIONS, 1962

TABLE 4.—FISSION PRODUCT GROSS BETA ACTIVITY IN SURFACE AIR, NASN, SECOND QUARTER 1962

[Concentrations in pc/m³]

| Station location | | No. of samples | Minimum | Maximum | Average | Station location | | No. of samples | Minimum | Maximum | Average |
|-----------------------|-------------------|----------------|---------|---------|---------|------------------|-------------------|----------------|---------|---------|---------|
| Alabama: | Birmingham | 6 | 4.1 | 15.7 | 9.7 | Michigan: | Detroit | 7 | 4.3 | 7.1 | 5.4 |
| | Mobile | 6 | 2.2 | 14.6 | 7.9 | | Grand Rapids | 4 | 4.4 | 6.2 | 5.1 |
| Alaska: | Anchorage | 7 | 1.4 | 6.6 | 3.7 | | Kalamazoo | 6 | 4.1 | 17.2 | 7.8 |
| | Pt. Woronzof * | 7 | 0.5 | 4.9 | 2.6 | | Lansing | 6 | 2.7 | 7.2 | 5.1 |
| Arizona: | Grand Canyon Pk * | 6 | 6.1 | 17.0 | 11.9 | | Saginaw | 6 | 3.8 | 11.4 | 8.0 |
| | Maricopa Co * | 6 | 6.3 | 33.8 | 15.2 | Minnesota: | Minneapolis | 6 | 2.2 | 7.2 | 4.7 |
| | Phoenix | 6 | 7.7 | 16.3 | 11.8 | | St. Paul | 5 | 3.4 | 6.7 | 5.3 |
| | Tucson | 6 | 6.0 | 18.2 | 11.0 | Mississippi: | Jackson | 5 | 2.9 | 9.5 | 6.9 |
| Arkansas: | Little Rock | 6 | 2.5 | 12.9 | 8.7 | | Jackson Co * | 5 | 1.5 | 10.0 | 6.5 |
| | Montgomery Co * | 5 | 3.2 | 12.5 | 7.5 | Missouri: | Kansas City | 5 | 5.0 | 9.8 | 6.7 |
| California: | Burbank | 7 | 2.5 | 20.2 | 8.5 | | Shannon Co * | 6 | 5.0 | 13.6 | 9.7 |
| | Fresno | 6 | 3.4 | 15.1 | 7.9 | | St. Louis | 6 | 2.9 | 10.3 | 6.8 |
| | Humboldt Co * | 6 | 0.8 | 3.5 | 1.9 | Montana: | Glacier Nat. Pk * | 6 | 2.7 | 6.9 | 5.2 |
| | Los Angeles | 7 | 1.7 | 20.1 | 6.5 | | Helena | 6 | 6.0 | 12.0 | 8.7 |
| | Oakland | 5 | 0.7 | 14.5 | 5.4 | Nebraska: | Lincoln | 7 | 3.1 | 10.9 | 6.9 |
| | Pasadena | 5 | 2.1 | 10.0 | 4.7 | | Omaha | 6 | 3.9 | 12.1 | 6.9 |
| | Richmond | 7 | 0.7 | 10.5 | 3.4 | | Thomas Co * | 7 | 5.1 | 16.3 | 8.0 |
| | Sacramento | 6 | 2.3 | 18.6 | 7.0 | Nevada: | Las Vegas | 5 | 6.1 | 23.6 | 14.7 |
| | San Diego | 7 | 2.3 | 12.3 | 6.5 | | White Pine Co * | 6 | 6.4 | 13.8 | 9.2 |
| | San Francisco | 6 | 0.6 | 10.9 | 3.9 | New Hamp- | | | | | |
| | Santa Barbara | 6 | 1.4 | 16.7 | 4.8 | shire: | Coos Co * | 7 | 3.1 | 8.6 | 6.4 |
| Colorado: | Denver | 6 | 4.4 | 7.3 | 5.5 | | Manchester | 5 | 4.4 | 8.9 | 6.2 |
| | Montezuma Co * | 5 | 5.0 | 15.0 | 8.6 | New Jersey: | Camden | 6 | 1.8 | 9.2 | 6.0 |
| Connecticut: | Bridgeport | 6 | 3.6 | 8.9 | 6.3 | | East Orange | 6 | 0.6 | 8.7 | 5.6 |
| | Hartford | 6 | 2.9 | 7.0 | 5.5 | | Elizabeth | 7 | 1.4 | 10.9 | 5.7 |
| | New Haven | 6 | 2.2 | 11.8 | 7.3 | | Newark | 6 | 1.5 | 7.2 | 4.5 |
| | Stamford | 6 | 2.1 | 11.7 | 5.6 | | Trenton | 6 | 1.7 | 9.0 | 5.6 |
| Delaware: | Kent Co * | 4 | 4.6 | 28.4 | 11.1 | New Mexico: | Albuquerque | 7 | 5.5 | 15.4 | 10.0 |
| | Wilmington | 7 | 1.9 | 8.0 | 5.0 | | Colfax Co * | 6 | 6.0 | 17.0 | 10.1 |
| District of Columbia: | Washington | 6 | 3.3 | 8.2 | 6.3 | New York: | Cape Vincent * | 5 | 6.3 | 9.1 | 7.6 |
| Florida: | Florida Keys * | 5 | 1.5 | 8.6 | 5.2 | | Elmira | 6 | 4.0 | 9.0 | 6.3 |
| | Jacksonville | 6 | 1.2 | 16.0 | 7.1 | | Glen Cove | 2 | 6.6 | 8.6 | 7.6 |
| | St. Petersburg | 7 | 1.9 | 13.0 | 7.1 | | Massena | 6 | 1.2 | 8.2 | 4.4 |
| | Tampa | 6 | 1.0 | 12.8 | 7.5 | | Mt. Vernon | 5 | 6.2 | 15.7 | 9.2 |
| Georgia: | Atlanta | 6 | 3.1 | 11.1 | 6.1 | | New Rochelle | 6 | 1.9 | 8.7 | 5.6 |
| | Columbus | 6 | 2.8 | 18.0 | 7.3 | | New York | 7 | 2.9 | 10.2 | 7.6 |
| | Macon | 6 | 2.8 | 9.6 | 6.8 | | Rochester | 6 | 1.0 | 9.2 | 5.8 |
| Hawaii: | Honolulu | 6 | 2.0 | 4.6 | 3.1 | | Troy | 4 | 1.1 | 9.8 | 6.4 |
| | Kahalaui * | 6 | 1.3 | 5.9 | 3.8 | North Carolina: | Asheville | 5 | 3.0 | 12.6 | 6.7 |
| Idaho: | Boise Idaho | 6 | 5.2 | 19.0 | 9.3 | | Cape Hatteras * | 6 | 6.0 | 14.1 | 9.5 |
| | Butte Co * | 7 | 5.5 | 11.0 | 8.6 | | Charlotte | 4 | 5.7 | 10.7 | 8.9 |
| Illinois: | Chicago | 7 | 3.2 | 10.3 | 6.5 | | Winston Salem | 6 | 2.5 | 19.1 | 9.6 |
| | Cicero | 6 | 2.1 | 7.0 | 5.2 | North Dakota: | Bismarek | 6 | 1.9 | 11.0 | 5.4 |
| | Peoria | 6 | 5.4 | 10.7 | 7.9 | | Ward Co * | 6 | 2.2 | 7.2 | 5.1 |
| | Springfield | 6 | 3.5 | 16.3 | 8.9 | Ohio: | Akron | 7 | 2.6 | 12.5 | 6.6 |
| Indiana: | East Chicago | 5 | 1.6 | 11.2 | 4.9 | | Cincinnati | 5 | 2.1 | 14.8 | 7.0 |
| | Evansville | 6 | 3.3 | 6.2 | 4.8 | | Cleveland | 6 | 4.7 | 10.0 | 7.6 |
| | Fort Wayne | 7 | 4.8 | 15.8 | 8.5 | | Columbus | 6 | 2.2 | 15.7 | 7.3 |
| | Gary | 6 | 4.9 | 12.8 | 7.4 | | Dayton | 6 | 2.2 | 13.1 | 6.5 |
| | Indianapolis | 6 | 4.0 | 11.6 | 7.4 | | Lorain | 7 | 2.8 | 11.0 | 6.9 |
| | Montgomery Co * | 6 | 3.6 | 11.4 | 7.5 | | Springfield | 7 | 3.9 | 11.4 | 7.2 |
| Iowa: | Clayton Co * | 6 | 2.6 | 16.1 | 7.5 | | Toledo | 6 | 5.7 | 10.0 | 7.2 |
| | Des Moines | 7 | 3.2 | 13.1 | 6.5 | | Youngstown | 7 | 4.6 | 10.3 | 7.6 |
| Kansas: | Kansas City | 6 | 1.9 | 10.5 | 5.9 | Oklahoma: | Cherokee Co * | 6 | 6.0 | 15.4 | 10.1 |
| | Wichita | 6 | 5.1 | 16.2 | 10.1 | | Oklahoma City | 7 | 7.6 | 13.4 | 10.1 |
| Kentucky: | Louisville | 5 | 3.1 | 9.0 | 5.6 | | Tulsa | 5 | 5.6 | 9.0 | 7.6 |
| Louisiana: | Baton Rouge | 7 | 2.2 | 14.4 | 9.6 | Oregon: | Curry Co * | 6 | 0.5 | 4.7 | 2.0 |
| | Lake Charles | 6 | 0.6 | 10.3 | 6.3 | | Portland | 5 | 0.9 | 12.8 | 5.1 |
| | New Orleans | 6 | 2.9 | 9.4 | 6.7 | Pennsyl- | | | | | |
| Maine: | Acadia Nat Pk * | 4 | 3.2 | 8.9 | 5.4 | vania: | Clarion Co * | 5 | 4.6 | 16.1 | 7.6 |
| | Portland | 6 | 2.9 | 7.1 | 4.9 | | Harrisburg | 5 | 0.7 | 10.0 | 6.6 |
| Maryland: | Baltimore | 6 | 5.6 | 16.7 | 8.5 | | Lancaster | 5 | 2.7 | 27.0 | 10.6 |
| | Calvert Co * | 6 | 5.3 | 11.6 | 8.0 | | Philadelphia | 7 | 2.4 | 9.5 | 6.0 |
| Massachu- | | | | | | | Pittsburgh | 6 | 1.9 | 11.4 | 6.7 |
| setts: | Boston | 7 | 1.3 | 11.5 | 6.4 | | Reading | 6 | 4.7 | 8.1 | 6.4 |
| | Fall River | 7 | 3.2 | 15.5 | 9.7 | | Wilkes Barre | 6 | 3.0 | 10.9 | 7.2 |
| | Lawrence | 6 | 2.6 | 11.7 | 7.7 | Puerto Rico: | Loquillo Mtns * | 5 | 2.1 | 9.9 | 4.3 |
| | Lynn | 6 | 3.7 | 10.4 | 6.1 | | San Juan | 5 | 2.0 | 7.8 | 4.1 |
| | Quincy | 5 | 5.5 | 10.3 | 8.5 | Rhode Island: | Providence | 6 | 6.7 | 10.7 | 8.0 |
| | Somerville | 6 | 4.0 | 12.3 | 7.4 | | Washington Co * | 7 | 0.5 | 6.9 | 4.9 |
| | Springfield | 6 | 1.0 | 7.4 | 5.2 | South Carolina: | Columbia | 4 | 6.6 | 12.3 | 9.5 |
| | | | | | | | Greenville | 7 | 4.1 | 12.7 | 9.2 |
| | | | | | | | Richland Co * | 6 | 3.1 | 12.6 | 7.3 |

TABLE 4.—FISSION PRODUCT GROSS BETA ACTIVITY IN SURFACE AIR, NASN, SECOND QUARTER 1962—Con.

[Concentrations in pc/m³]

| Station location | | | | | Station location | | | | |
|----------------------------------|--|--|--|--|---------------------------|--|--|--|--|
| No. of samples | | | | | No. of samples | | | | |
| Minimum | | | | | Minimum | | | | |
| Maximum | | | | | Maximum | | | | |
| Average | | | | | Average | | | | |
| South Dakota: Black Hills Frst * | | | | | Virginia: Danville | | | | |
| Sioux Falls | | | | | Norfolk | | | | |
| Tennessee: Chattanooga | | | | | Shenandoah Nat. Pk * | | | | |
| Memphis | | | | | Washington: Clallam Co * | | | | |
| Nashville | | | | | Seattle | | | | |
| Texas: Aransas Co * | | | | | Tacoma | | | | |
| Corpus Christi | | | | | West Virginia: Charleston | | | | |
| Dallas | | | | | Huntington | | | | |
| El Paso | | | | | Wisconsin: Door Co * | | | | |
| Ft. Worth | | | | | Milwaukee | | | | |
| Houston | | | | | Racine | | | | |
| San Antonio | | | | | Wyoming: Cheyenne | | | | |
| Waco | | | | | Yellowstone Pk * | | | | |
| Utah: Salt Lake City | | | | | | | | | |
| Vermont: Burlington | | | | | | | | | |
| Orange Co * | | | | | | | | | |

*Nonurban station

Gross Beta Activity In Precipitation

During 1959 a precipitation collection and analysis program was established by the Weather Bureau Research Station in Cincinnati, Ohio, and the National Air Sampling Network. The collection stations are located at Weather Bureau offices or airport stations. Monthly composite samples of precipitation are

collected at 29 stations and forwarded to the Network laboratory for analysis.

Samples are analyzed for total solids and a large number of metals and nonmetals. In addition, samples are analyzed for fission product gross beta radioactivity if a large enough volume remains after the requirements for the chemical analysis have been met. Second quarter 1962 data are presented in table 5.

TABLE 5.—FISSION PRODUCT GROSS BETA ACTIVITY IN PRECIPITATION, NASN, SECOND QUARTER 1962

| Station | | April | | May | | June | |
|-----------------------|-----------------------|----------|-------------------|----------|-------------------|----------|-------------------|
| | | pc/liter | nc/m ² | pc/liter | nc/m ² | pc/liter | nc/m ² |
| Alabama | Montgomery | 910 | 140 | — | — | 890 | 111 |
| Florida | Tampa | * | — | 1030 | 66 | 439 | 32 |
| Illinois | Chicago, (Midway) | — | — | 1910 | 83 | 1170 | 65 |
| Illinois | Chicago, (O'Hare) | — | — | 2280 | 168 | — | — |
| Louisiana | Lake Charles | — | — | — | — | 504 | 66 |
| Maine | Caribou | 2010 | 85 | 1490 | 70 | 1180 | 60 |
| Massachusetts | Nantucket | 1190 | 107 | — | — | 474 | 43 |
| Michigan | Sault Ste. Marie | 1470 | 64 | 875 | 105 | — | — |
| Minnesota | St. Cloud | — | — | 1480 | 130 | 1830 | 115 |
| Montana | Glasgow | — | — | 1140 | 66 | 1160 | 76 |
| Nebraska | Grand Island | — | — | 1980 | 167 | 843 | 53 |
| New York | Albany | 1130 | 48 | 2010 | 95 | — | — |
| North Carolina | Cape Hatteras | 795 | 93 | — | — | 291 | 64 |
| Ohio | Cincinnati (Airport) | — | — | 1270 | 101 | 876 | 65 |
| | Cincinnati (Gest St.) | — | — | 1550 | 77 | 1010 | 87 |
| Pennsylvania | Philadelphia | 1300 | 97 | 2330 | 101 | 459 | 80 |
| South Carolina | Charleston | 2300 | 100 | — | — | 271 | 92 |
| South Carolina | Greenville | 1030 | 144 | — | — | 546 | 96 |
| Tennessee | Nashville | 785 | 114 | — | — | 643 | 107 |
| Texas | Amarillo | — | — | — | — | 1880 | 288 |
| Texas | Brownsville | — | — | 967 | 42 | 265 | 21 |
| Texas | San Angelo | 2460 | 148 | — | — | — | — |
| Virginia (Wash. D.C.) | Sterling | 2700 | 124 | 1750 | 76 | 521 | 45 |
| Washington | Tatoosh | 870 | 157 | — | — | — | — |

* Dash denotes no sample.

CANADIAN RADIOACTIVE FALLOUT
STUDY PROGRAM
First Quarter 1962

Department of National Health and Welfare,
Ottawa, Canada

As part of its Radioactive Fallout Study Program (RFSP), the Radiation Protection Division, Department of National Health and Welfare, Dominion of Canada, conducts air and precipitation sampling programs. The 24 air

and precipitation stations are located at airports, and the equipment is operated by meteorologists of the Meteorological Services Branch of the Department of Transport.

Air

About 650 cubic meters of air are drawn through a high efficiency 4-inch diameter filter during a 24-hour period. Filters are sent daily to the Radiation Protection Division Laboratory in Ottawa. A 2-inch diameter disk is removed from each filter and counted with a thin end-window Geiger flow counter system, cali-

TABLE 6.—FISSION PRODUCT GROSS BETA ACTIVITY IN SURFACE AIR, RFSP, FIRST QUARTER 1962

[Average concentration in $\mu\text{mc}/\text{m}^3$]

| Station location | January | February | March | Station location | January | February | March |
|--------------------|---------|----------|-------|--------------------------|---------|----------|-------|
| Calgary..... | 8.9 | 7.6 | 9.0 | Resolute..... | 2.2 | *— | 9.6 |
| Chatham..... | 8.4 | 8.9 | 6.2 | Saskatoon..... | 7.9 | 6.3 | 6.0 |
| Coral Harbour..... | 6.0 | 5.6 | 5.2 | Shearwater..... | 10.3 | 8.9 | 6.2 |
| Edmonton..... | 9.7 | 7.8 | 6.4 | Toronto..... | 7.8 | 4.1 | — |
| Ft. Churchill..... | 4.8 | 5.1 | 5.6 | Toronto..... | 9.4 | 9.3 | 8.4 |
| Ft. William..... | 8.9 | 7.9 | 8.0 | Vancouver..... | 6.1 | 10.9 | 6.6 |
| Fredericton..... | 8.7 | 9.8 | 6.0 | Whitehorse..... | 6.5 | 9.4 | 7.9 |
| Goose Bay..... | 5.9 | 6.7 | 4.7 | Windsor..... | 9.3 | 8.8 | 9.5 |
| Inuvik..... | 8.0 | 7.8 | 6.2 | Winnipeg..... | 8.6 | 7.2 | 7.4 |
| Kapuskasing..... | 8.2 | 8.4 | 9.7 | Yellowknife..... | 7.3 | 6.2 | 6.9 |
| Montreal..... | 8.5 | 7.2 | 7.5 | Maximum station ave..... | 10.3 | 10.9 | 9.7 |
| Moosonee..... | 8.2 | 7.7 | 9.4 | Minimum station ave..... | 2.2 | 4.1 | 4.7 |
| Ottawa..... | 9.2 | 7.7 | 8.5 | Overall average..... | 7.8 | 7.6 | 7.3 |
| Regina..... | 8.7 | 6.7 | 6.9 | | | | |

* Dash denotes no sample.

TABLE 7.—FISSION PRODUCT GROSS BETA ACTIVITY IN PRECIPITATION, RFSP, FIRST QUARTER 1962

| Station location | January | | February | | March | |
|--------------------|----------------------------------------------|----------------------------------------------------|----------------------------------------------|----------------------------------------------------|----------------------------------------------|----------------------------------------------------|
| | Concentration $\mu\text{mc}/\text{liter}$ | Deposition ($\text{m}\mu\text{c}/\text{m}^2$) | Concentration $\mu\text{mc}/\text{liter}$ | Deposition ($\text{m}\mu\text{c}/\text{m}^2$) | Concentration $\mu\text{mc}/\text{liter}$ | Deposition ($\text{m}\mu\text{c}/\text{m}^2$) |
| Calgary..... | 1030 | 12.1 | 467 | 5.2 | 1460 | 15.2 |
| Chatham..... | 751 | 48.8 | 2040 | 48.8 | 751 | 55.0 |
| Coral Harbour..... | a tr | 14.5 | tr | 7.7 | 2430 | 21.0 |
| Edmonton..... | 1560 | 39.2 | 923 | 35.8 | 1340 | 34.1 |
| Fredericton..... | 739 | 47.1 | 812 | 56.3 | 2830 | 77.5 |
| Ft. Churchill..... | 70 | 12.5 | 1310 | 5.7 | — | — |
| Ft. William..... | 2240 | 33.0 | 885 | 50.7 | 2700 | 40.5 |
| Goose Bay..... | 328 | 16.2 | 391 | 19.7 | 874 | 64.7 |
| Inuvik..... | 675 | 20.9 | 873 | 21.9 | 918 | 11.0 |
| Kapuskasing..... | 458 | 18.3 | 272 | 7.9 | 1120 | 24.2 |
| Montreal..... | 3020 | 118.8 | 1490 | 114.4 | 1260 | 84.6 |
| Moosonee..... | b— | — | — | — | 806 | 28.0 |
| Ottawa..... | 1100 | 62.6 | 638 | 41.1 | 1080 | 50.6 |
| Regina..... | 2510 | 45.2 | 2420 | 31.3 | 1690 | 32.3 |
| Resolute..... | tr | 52.0 | tr | 96.7 | tr | 56.0 |
| Saskatoon..... | 2990 | 45.5 | 999 | 36.8 | 876 | 16.9 |
| Shearwater..... | — | — | 1090 | 171.5 | 1090 | 57.9 |
| Toronto..... | 505 | 53.1 | 648 | 140.0 | 1020 | 150.0 |
| Toronto..... | 1950 | 107.8 | 1520 | 113.3 | 2870 | 29.1 |
| Vancouver..... | — | — | 565 | 36.1 | 768 | 71.8 |
| Whitehorse..... | 927 | 24.7 | 2380 | 23.5 | 789 | 18.6 |
| Windsor..... | 1590 | 87.7 | 1690 | 133.1 | 1420 | 36.7 |
| Winnipeg..... | 1550 | 34.3 | 793 | 31.6 | 1380 | 41.6 |
| Yellowknife..... | 602 | 17.3 | 471 | 6.0 | 128 | 8.4 |
| Average..... | 1290 | 43.4 | 1080 | 53.7 | 1340 | 44.6 |

a tr denotes trace precipitation.

b Dash denotes no sample.




FIGURE 6.—CANADIAN AIR AND PRECIPITATION SAMPLING STATIONS

brated with a $\text{Sr}^{90}\text{-Y}^{90}$ standard. The delay period between collection and counting is one week to avoid interference from naturally-occurring radon and thoron daughters. Air data for first quarter 1962 are presented in table 6.

Precipitation

The amount of radioactive fallout being deposited on the ground is determined from meas-

urements on material collected in special polythene-lined rainfall pots. After transfer of the water to the sampling container, the polythene liner is removed, packed with the sample, and sent to the laboratory. First quarter precipitation data are presented in table 7.

More detailed discussions of the sampling procedures, methods of analysis, and interpretation of results of the radioactive fallout program are contained in the Department's annual reports for 1959 and 1960 (1, 2).

REFERENCES

- (1) Bird, P. M., A. H. Booth, and P. G. Mar: *Annual Report for 1959 on the Radioactive Fallout Study Program, CNHW (RP-3)*, (May 1960).
- (2) Bird, P. M., A. H. Booth, and P. G. Mar: *Annual Report for 1960 on the Radioactive Fallout Study Program, CNHW (RP-4)*, (December 1961).

Previous coverage in *Radiological Health Data*:

| Period | Issue |
|---------------------|----------------|
| July 1959-June 1960 | January 1961 |
| Third quarter 1960 | April 1961 |
| Fourth quarter 1960 | August 1961 |
| First quarter 1961 | December 1961 |
| Second quarter 1961 | February 1962 |
| Third quarter 1961 | May 1962 |
| Fourth quarter 1961 | September 1962 |

Natural and Fission Product Radioactivity in Surface Air Particulates at Cincinnati, Ohio

June 18-July 13, 1962


Division of Radiological Health, Public Health Service

The determination of natural radioactivity in the atmosphere is useful because the exposure levels from natural radiation can be used as a base for comparative evaluations of exposures from artificially produced radionuclides. Some

of the more important examples of naturally occurring and artificially produced airborne radioactivity are given in table 1. The physical state is of importance in the selection of sampling procedure.

TABLE 1.—EXAMPLES OF AIRBORNE RADIOACTIVITY

| Physical state | Natural | | Artificial |
|----------------|---------------------------------------------------------------------------------------------|-------------------------|----------------------------------------------|
| | Nuclides (radiation) | α half life | |
| Gaseous | Radon (α, γ) Thoron (α, γ) | 3.8 days 54 seconds | Some fission products (β, γ) |
| Particulates | Radon daughters (α, β, γ) Thoron daughters (α, β, γ) | 0.5 hours 10.6 hours | Most fission products (β, γ) |



* Equilibrium values do not in fact remain constant throughout a 24-hour period but vary with changes of radon and thoron concentrations in air.

FIGURE 1.—IDEALIZED PROFILE OF ALPHA ACTIVITY ON FILTER DURING AND AFTER SAMPLING

Radon and Thoron

The earth's crust contains trace amounts of naturally-occurring uranium and thorium, each of which decays through a series of radionuclides, ending finally in stable isotopes of lead. One of the radionuclides in the uranium series is the inert gas, radon-222, commonly called "radon." In the thorium series, the inert gas decay product is radon-220, or "thoron." Some radon and thoron gases formed near the ground surface escape into the atmosphere before decaying to their particulate daughter products. Concentrations of radon and thoron in surface air depend on prevailing atmospheric conditions such as ambient temperature, wind, and pressure and on soil conditions such as moisture, porosity, and temperature that affect the rate of diffusion of the gases through the soil.

Most of the natural radioactivity in surface

air is due to radon and its daughters. Thoron and its daughters contribute less because of thoron's shorter half life, and therefore, less time for diffusion through the pores of the soil to escape into the atmosphere. The natural radioactivity concentration in air shows a pronounced daily fluctuation. During a period of atmospheric inversion, such as often occurs in the early morning, radon and thoron tend to accumulate in the air near the ground surface. During periods of atmospheric turbulence, radon and thoron are dispersed in the atmosphere, leaving lower concentrations near the surface.

Measurement of radon and thoron concentrations in air may be accomplished by two general methods. The direct method involves the collection of radon and thoron gases by charcoal adsorption (1) while the indirect method utilizes the collection of airborne particulates,

TABLE 2.—SURFACE AIR RADON (Rn^{222}), THORON (Rn^{220}), AND FISSION PRODUCT GROSS BETA CONCENTRATIONS AT CINCINNATI, OHIO, JUNE 18–JULY 13, 1962

| End of sampling period | | Rn^{222} 8 a.m. ($\mu\mu\text{C}/\text{m}^3$) | Rn^{222} 3 p.m. ($\mu\mu\text{C}/\text{m}^3$) | Rn^{220} ($\mu\mu\text{C}/\text{m}^3$) | Beta activity ($\mu\mu\text{C}/\text{m}^3$) |
|----------------------------------------------|---------|----------------------------------------------------------------|----------------------------------------------------------------|------------------------------------------------------|-----------------------------------------------------|
| June 18 | ----- | 880 | 160 | 3.0 | 9.53 |
| 19 | ----- | 240 | 150 | 1.8 | 6.89 |
| 20 | ----- | 460 | 140 | 4.8 | 7.21 |
| 21 | ----- | 620 | 130 | 6.8 | 5.55 |
| 22 | ----- | 440 | 200 | 5.4 | 8.65 |
| 25 | ----- | 700 | 160 | 4.2 | 5.63 |
| 26 | ----- | 770 | 160 | 6.7 | 9.10 |
| 27 | ----- | 410 | 240 | 4.7 | 7.80 |
| 28 | ----- | 240 | 110 | 9.6 | 7.16 |
| 29 | ----- | 610 | 140 | 6.5 | 7.93 |
| July 2 | ----- | 600 | 190 | 6.7 | 7.48 |
| 3 | ----- | 260 | 200 | 2.9 | 5.72 |
| 5 | ----- | 480 | 200 | 3.3 | 1.91 |
| 6 | ----- | 500 | 190 | 3.5 | 1.41 |
| 9 | ----- | 250 | 70 | 3.5 | 2.98 |
| 10 | ----- | 690 | 120 | 5.4 | 3.24 |
| 11 | ----- | 670 | 160 | 6.7 | 5.21 |
| 12 | ----- | 250 | 130 | 3.1 | 4.87 |
| 13 | ----- | 500 | 170 | 5.4 | 4.94 |
| Average | | 532 | 156 | 4.7 | 5.94 |
| Range of counting errors (2 σ) | Maximum | 52 | 27 | 1.0 | 0.18 |
| | Minimum | 27 | 15 | 0.4 | 0.06 |

which include the radon and thoron daughter products, on the surface of a membrane filter. The choice of the indirect method for routine monitoring was made because of simplicity, sensitivity, and low cost.

The measurement of radon and thoron by the indirect method is based on the assumption that radon and thoron in the atmosphere are in equilibrium with their daughters. Under these conditions the particulate alpha activity (from the daughter products) retained on the membrane filter by a uniform flow of air may be theoretically related to the concentrations of radon and thoron in air. If radon and thoron activities in the air remain constant, the alpha activity on the filter while the sampler is in operation will reach 97 percent of equilibrium in 2.5 hours for radon daughters and in 53 hours for thoron daughters. The measurement of the activity by alpha scintillation, which is specific for alpha activity, eliminates interference from fission product beta activity.

The method used for the separate measurements of radon and thoron daughters is diagrammed in figure 1 where the history of one filter during a typical 24-hour sampling period

and for a period following its removal is charted. At 3:00 p.m. the filter is removed momentarily from the sampler and counted. Although some thoron daughter activity is present, it is neglected and the activity is taken as radon daughter activity. The time of 3:00 p.m. was selected to represent the time of day most likely to have atmospheric turbulence and, therefore, lowest radon concentration in surface air. Radon and thoron concentrations often reach a maximum during an early morning atmospheric inversion condition. The 8:00 a.m. measurement of activity therefore approximates the daily maximum.

The measured alpha activities are adjusted to 100 percent equilibrium with factors based on length of sampling period and decay time after removal of filter. (2) The equilibrium alpha activities on the filter due to the respective daughter products may in turn be converted to radon and thoron concentrations in air by the following expressions:

$$\text{radon } (\mu\mu\text{C}/\text{m}^3) = 6 A_r/F, \text{ and}$$

$$\text{thoron } (\mu\mu\text{C}/\text{m}^3) = 0.45 A_t/F,$$

where A_r and A_t are the radon and thoron daughter alpha activities (dpm) respectively and F is the air sampling rate in liters per minute at the end of the sampling period.

Table 2 shows daily calculated radon activities for 8 a.m. and 3 p.m. and thoron for 8 a.m.

Fission Product Activity

The membrane filter used for the determination of radon and thoron is retained several days to allow for the decay of thoron daughters. At that time the beta activity is measured. This activity represents fission products and is reported in table 2.

REFERENCES

- (1) Shleien, B.: The Simultaneous Determination of Atmospheric Radon by Filter Paper and Charcoal Adsorptive Techniques, Paper presented before 23rd Meeting of the American Industrial Hygiene Association and submitted for publication in the *American Industrial Hygiene Association Journal* on May 17, 1962.
- (2) Setter, L. R., and G. I. Coats: The Determination of Airborne Radioactivity, *American Industrial Hygiene Association Journal*, 22: 64-9 (Feb. 1962).

SECTION II.—FOOD

Strontium-90 Concentrations in U.S. Wheat and Milling Products¹

1960

J. Rivera

Health and Safety Laboratory, U.S. Atomic Energy Commission

Concentrations of strontium-90 and calcium in U.S. wheat and milling products have been studied extensively at the Health and Safety Laboratory (HASL) since 1958. Table 2 compares the results of the survey of the 1960 crop with those previously published for the 1958 and 1959 crops. A pictorial representation of the data for 1959 and 1960 is presented in figure 1. Each cube, scaled to represent the concentration, is placed on the appropriate state. Material balances for the milling products are given in table 3, and new information on the 1960 milling products² is summarized in table

4. These data show a remarkable similarity in the relative distributions of strontium-90 among the different milling fractions. Moreover, this distribution pattern appears to have persisted over the three year period, in spite of the fact that fallout conditions varied considerably. The logical conclusion is that most

TABLE 1.—STRONTIUM-90 DEPOSITION IN WHEAT AND SOIL

| Year | Wheat ($\mu\mu\text{c Sr}^{90}/\text{g Ca}$) | Soil Description | |
|-----------|---------------------------------------------------|------------------------------------------------------------|---------------------------------------------------------|
| | | Cumulative ($\text{m}\mu\text{c Sr}^{90}/\text{m}^2$) | ($\text{m}\mu\text{c Sr}^{90}/\text{m}^2$) in June |
| 1958..... | 172* | 18 | 0.58 |
| 1959..... | 147* | 26 | 0.57 |
| 1960..... | 62 | 27 | 0.20 |

* Assuming 0.36 g Ca/kg of wheat.

¹ U.S. Atomic Energy Commission: *Summary Quarterly Reports, HASL-122*, Office of Technical Services, Department of Commerce, Washington 25, D.C. (April 1, 1962), price \$3.00.

² The production of flour from wheat grain is accomplished through milling. The first stage in this process involves the flattening and splitting of the kernels and the removal of the wheat *germ* which is not utilized in the production of flour. The grain is then passed through a series of rollers which remove successive layers of cells from the kernel. The highest grade flour is made mostly from the innermost part of the kernel and is called *patent flour*. Flour of lower quality is made from layers of cells nearer to the surface of the kernel, this flour is called the *first or second clear*. The outermost part of the wheat berry which forms a coat which is relatively easily removed is called the *bran*. Bran is not used in making flour. *Shorts* consists mainly of fine particles of bran mixed with some clear flour.

of the strontium-90 found in wheat originated from a single source.

Strontium-90 in wheat is derived either from the soil by root extraction or through direct deposition on foliar surfaces. Minimum estimates of strontium-90 in wheat from the latter source have been made by comparing the specific activity ($\mu\mu\text{c Sr}^{90}/\text{mg}$ stable Sr) of flour with that of bran and assuming that the strontium-90 in the flour is derived solely from

TABLE 2.—COMPARISON BETWEEN STRONTIUM-90 AND CALCIUM LEVELS IN UNITED STATES WHEAT SAMPLED IN 1958, 1959, AND 1960

| State | Calcium | | | Strontium-90 | | | | | |
|-------------------|---------|------|------|--------------|------|------|------------|---------------------|--------|
| | (g/kg) | | | (μμc/kg) | | | (μμc/g Ca) | | |
| | 1958 | 1959 | 1960 | 1958 | 1959 | 1960 | 1958 | 1959 | 1960 |
| California..... | — | 0.34 | 0.33 | — | 17 | 3.7 | — | 49 (2) ^b | 11 (5) |
| Colorado..... | — | 0.49 | 0.42 | — | 46 | 19 | — | 71 (5) | 46 (2) |
| Idaho..... | — | 0.34 | 0.35 | — | 5.7 | 6.1 | — | 17 (1) | 18 (2) |
| Illinois..... | 0.40 | 0.38 | — | 133 | 54 | — | 330 (1) | 142 (1) | — |
| Iowa..... | — | 0.54 | 0.38 | — | 62 | 34 | — | 109 (2) | 89 (1) |
| Kansas..... | 0.36 | 0.49 | 0.42 | 82 | 47 | 26 | 227 (1) | 97 (1) | 62 (1) |
| Michigan..... | 0.36 | 0.38 | — | 42 | 48 | — | 118 (1) | 125 (1) | — |
| Minnesota..... | 0.35 | 0.33 | — | 51 | 59 | — | 162 (16) | 181 (2) | — |
| Missouri..... | — | 0.43 | 0.42 | — | 121 | 21 | — | 272 (4) | 50 (1) |
| Montana..... | 0.47 | 0.38 | 0.40 | 59 | 20 | 15 | 125 (1) | 52 (1) | 38 (1) |
| Nebraska..... | — | 0.48 | 0.37 | — | 76 | 32 | — | 159 (3) | 68 (3) |
| New York..... | 0.36 | 0.37 | 0.41 | 44 | 44 | 16 | 121 (1) | 119 (1) | 40 (1) |
| North Dakota..... | 0.31 | 0.33 | 0.29 | 40 | 51 | 26 | 134 (10) | 154 (1) | 91 (1) |
| Ohio..... | — | 0.36 | 0.48 | — | 56 | 35 | — | 152 (11) | 73 (4) |
| Oklahoma..... | 0.41 | 0.40 | 0.39 | 34 | 63 | 35 | 82 (1) | 159 (1) | 90 (2) |
| South Dakota..... | 0.35 | — | — | 61 | — | — | 171 (8) | — | — |
| Texas..... | 0.44 | 0.51 | — | 44 | 51 | — | 99 (2) | 99 (1) | — |
| Utah..... | — | 0.48 | 0.44 | — | 21 | 7.9 | — | 45 (5) | 19 (4) |
| Washington..... | — | 0.32 | 0.35 | — | 9.2 | 7.0 | — | 30 (2) | 20 (1) |

^a Dash indicates no sample.

^b Numbers in parentheses indicate number of samples averaged.

the soil by root uptake (1). The inaccuracy of this assumption is apparent from the data presented in table 3. Despite the fact that the strontium-90 levels in soil probably increased in 1960 as compared to 1959, the strontium-90 concentration in patent flour decreased sharply. It follows that much of the strontium-90 in the patent flour was derived by translocation of strontium-90 directly deposited on the outer parts of the grain. This translocation effect

should be more easily seen in 1962 by tracing fresh fallout expected to be found in wheat through Sr^{89}/Sr^{90} ratios in the milling fractions. Presumably this will be possible since it is expected that the 1962 wheat crop will be contaminated by fresh fallout from the Soviet test series that were held in late 1961.

The concentration of strontium-90 in wheat grown in the United States was estimated to be $62 \pm 27 \mu\mu\text{c/kg}$ in 1958; $53 \pm 19 \mu\mu\text{c/kg}$ in

TABLE 3.—STRONTIUM-90 MATERIAL BALANCE IN UNITED STATES WHEAT AND MILLING PRODUCTS

| Material | Weight (kg) | ^a 1958 | | ^b 1959 | | ^c 1960 | |
|---------------------------|-------------|-------------------|-----------------|-------------------|-----------------|-------------------|-----------------|
| | | μμc/kg | μμc in fraction | μμc/kg | μμc in fraction | μμc/kg | μμc in fraction |
| Input: Wheat..... | 100 | 62 | 6200 | 53 | 5300 | 28 | 2800 |
| Output: Patent Flour..... | 58 | 12 | 696 | 9 | 522 | 4 | 232 |
| 1st & 2nd Clear..... | 14 | 28 | 392 | 17 | 238 | 9 | 126 |
| Germ & Shorts..... | 17 | 133 | 2261 | 143 | 2431 | 66 | 1122 |
| Bran..... | 11 | 231 | 2541 | 163 | 1793 | 107 | 1177 |
| Output total..... | 100 | | 5890 | | 4984 | | 2657 |
| Difference..... | | | 310 (5%) | | 316 (6%) | | 143 (5%) |

^a Production weighted average of 9 states.

^b Production weighted average of 5 states representing 64 percent of 1958 production, *The World Almanac and Book of Facts for 1960*.

^c Production weighted average of 6 states representing 55 percent of 1960 production, *The World Almanac and Book of Facts for 1962*.




FIGURE 1.—STRONTIUM-90 IN U.S. WHEAT HARVESTED IN 1959 & 1960

TABLE 4.—STRONTIUM-90 AND CALCIUM IN WHEAT AND MILLING PRODUCTS 1960

| State | Wheat | Bran | Germ | Shorts | Clear | Patent |
|----------------------------------|-------|------|------|--------|-------|--------|
| <i>μC Sr⁹⁰/kg:</i> | | | | | | |
| Kansas..... | 26 | 111 | 49 | 97 | 9.4 | 4.6 |
| Montana..... | 15 | 70 | 70 | 49 | 4.5 | 3.3 |
| Nebraska..... | 32 | 115 | 60 | 26 | 8.8 | 4.9 |
| North Dakota..... | 26 | 102 | 42 | 76 | 7.2 | 3.8 |
| Ohio..... | 31 | 95 | 37 | 56 | 12 | 5.4 |
| Oklahoma—(80%), Texas—(20%)..... | 44 | 138 | 61 | 110 | 11 | 3.8 |
| <i>μC Sr⁹⁰/g Ca:</i> | | | | | | |
| Kansas..... | 62 | 78 | 67 | 90 | 43 | 22 |
| Montana..... | 38 | 81 | 159 | 58 | 30 | 36 |
| Nebraska..... | 85 | 125 | 108 | 23 | 35 | 31 |
| North Dakota..... | 90 | 111 | 81 | 87 | 48 | 34 |
| Ohio..... | 78 | 68 | 61 | 80 | 46 | 36 |
| Oklahoma—(80%), Texas—(20%)..... | 113 | 141 | 95 | 119 | 38 | 20 |
| <i>g Ca/kg:</i> | | | | | | |
| Kansas..... | 0.42 | 1.4 | 0.74 | 1.1 | 0.22 | 0.21 |
| Montana..... | 0.40 | 0.86 | 0.42 | 0.87 | 0.15 | 0.09 |
| Nebraska..... | 0.37 | 0.92 | 0.56 | 1.1 | 0.25 | 0.16 |
| North Dakota..... | 0.29 | 0.93 | 0.52 | 0.87 | 0.15 | 0.11 |
| Ohio..... | 0.41 | 1.4 | 0.61 | 0.70 | 0.26 | 0.15 |
| Oklahoma—(80%), Texas—(20%)..... | 0.39 | 0.97 | 0.63 | 0.94 | 0.29 | 0.19 |

1959; and 24 ± 9 $\mu\text{C}/\text{kg}$ in 1960.³ While the difference between the 1958 and 1959 concentrations was not statistically significant, the difference between the 1959 and 1960 con-

³ Weighted average of 14 states representing 73 percent of 1960 production. *The World Almanac and Book of Facts for 1962*. The error terms given for strontium-90 concentrations in U.S. wheat are one standard deviation from the production weighted average.

centrations is significant. Since the level of strontium-90 in the soil increased between 1959 and 1960, the decline in the concentration of strontium-90 in wheat grown in 1960 must have been due to a decline in the fallout rate and a consequent decrease in direct deposition of strontium-90 on the grain.

Middleton has shown that wheat is most

likely to be contaminated by direct deposition after the emergence of the ears from the boot (2). For the majority of wheat harvested in the United States, this occurs in June (3). The fallout rate in June for the midwestern wheat producing states can be estimated from the HASL monthly pot and funnel monitoring network. The cumulative deposit of strontium-90 in the soils of these states can also be estimated from the HASL-USDA soil sampling program. These estimates, together with the Sr^{90}/Ca ratio in wheat for 1958, 1959 and 1960, are shown in table 1.

From these data in table 1 we can find suitable values for proportionality factors "a" and "b" in an expression of the form,

$$V = a S + b R$$

where V is the Sr^{90}/Ca ratio in the wheat, S is the cumulative deposit of strontium-90 in the soil, and R is the fallout rate. A choice of "a" and "b" that fit the data reasonably well is, $a = 0.12 \mu\text{c/g Ca/mc/mi}^2$ and $b = 102 \mu\text{c/g Ca/mc/mi}^2$ deposited in June. Using these

values for "a" and "b," the fraction of the Sr^{90}/Ca ratio in wheat due to direct deposition was about 90 percent for all three years. This result is in good agreement with the estimate of Menzel, et al. (4) for wheat grown in Maryland in 1959.

REFERENCES

- (1) Agricultural Research Council Radiobiological Laboratory: *Strontium-90 in Human Diet in the United Kingdom-1958*, ARCRL 1, Her Majesty's Stationery Office, London (1959), p. 47.
- (2) Middleton, L. J.: Radioactive Strontium and Cesium in the Edible Parts of Crop Plants after Foliar Contamination, *International Journal of Radiation Biology*, 1:387 (October 1959).
- (3) Private Communication, L. W. Briggie: U.S. Department of Agriculture, Beltsville, Maryland.
- (4) Menzel, R. G., et al: Foliar Retention of Strontium-90 by Wheat, *Science*, 134:559-60 (August 25, 1961).

Previous coverage of wheat data in *Radiological Health Data*:

| Period | Issue |
|----------------------|---------------|
| 1958-1959 | May 1960 |
| 1958-1960 | October 1960 |
| 1958-1959 | November 1960 |
| 1958-1959 | April 1961 |
| 1959 (foreign) | April 1961 |
| 1958-1959 (Canadian) | July 1961 |
| 1960 | January 1962 |

Deposition of Strontium-90 in a 1960 Kansas Wheat Sample¹

J. Rivera

Health and Safety Laboratory, U.S. Atomic Energy Commission

Knowledge of the mechanism by which plants are contaminated by fallout is essential for the accurate prediction of future levels of strontium-90 in foods. A particularly important fact that must be known is whether plants become contaminated principally from strontium-90 that is absorbed from the soil or from strontium-90 that results from the direct deposition of fallout on the plant surfaces. The experiment presented here on the contamination of wheat may be of some help in resolving this problem.

A sample of wheat from Kansas harvested in 1960 was treated in such a way as to remove the epidermal layer of the bran coats. The resulting two fractions, together with an aliquot of the original wheat sample, were ana-

lyzed for strontium-90 and stable strontium. The results are presented in table 1.

If all of the strontium-90 in the wheat berry were derived from the soil, then the specific activity ($\mu\text{c Sr}^{90}/\text{mg Sr}$) in each of the fractions of the berry would be the same. The outermost part of the berry, however, was exposed to direct deposition of strontium-90 from fallout in addition to that which was derived from the soil. If there had been no direct deposition of strontium-90 on the "beeswing," it would have had the same specific activity as the inner part of the berry, namely, $5.69 \mu\text{c Sr}^{90}/\text{mg Sr}$, and would have contained $5.69 \times 0.71 = 4.00 \mu\text{c Sr}^{90}$. Since the observed amount of strontium-90 in this fraction was $7.1 \mu\text{c}$, the difference of $3.1 \mu\text{c}$ could be attributed to direct deposition.

The strontium-90 in the stripped fraction of the wheat may not have originated entirely from strontium-90 in the soil as was assumed here, but rather some may have come from

¹ U.S. Atomic Energy Commission: *Summary Quarterly Report, HASL-127*, Office of Technical Services, Department of Commerce, Washington 25, D.C. (July 1, 1962), price \$3.00.

TABLE 1.—STRONTIUM IN WHEAT FRACTION

| Wheat fraction | $\mu\text{c Sr}^{90}/\text{kg}$ | kg in fraction | $\mu\text{c Sr}^{90}$ in fraction | mg Sr/kg | mg Sr in fraction | Specific activity ($\mu\text{c Sr}^{90}/\text{mg Sr}$) |
|---------------------|---------------------------------|----------------|-----------------------------------|----------|-------------------|----------------------------------------------------------|
| Wheat..... | 26.1 | 1.000 | 26.1 | 4.02 | 4.02 | 6.49 |
| Stripped wheat..... | 19.6 | 0.968 | 19.0 | 3.45 | 3.34 | 5.69 |
| Beeswing *..... | 216.4 | 0.033 | 7.1 | 21.57 | 0.71 | 10.00 |

* Fraction stripped from wheat.

strontium-90 that was directly deposited on the outer part of the berry and then translocated to the inner part. The estimate that $4.00 \mu\text{c Sr}^{90}$ in the "beeswing" came from the soil is, therefore, a maximum estimate and $3.1 \mu\text{c}$ must, therefore, be regarded as the minimum amount that was due to direct deposition.

There is considerable evidence to the effect that the amount of strontium-90 directly deposited on the wheat berry is proportional to the fallout rate (see page 392). One would expect, therefore, that the fraction of strontium-90 in wheat due to direct deposition would have been greater in a sample harvested in 1959 than one harvested in 1960 since the fallout was greater in 1959 than in 1960.

An experiment identical to the one described above was performed on a sample of Kansas wheat harvested in 1959 (1). The result was that an estimated 22 percent of the total strontium-90 in the wheat was the minimum amount that was ascribable to the direct deposition. The minimum amount of strontium-90 in the wheat due to direct deposition for the 1960 sample of $3.1 \mu\text{c}$ was 12 percent of the total in the sample. These results substantiate the belief that direct deposition constitutes a very important factor in the contamination of wheat with strontium-90 from fallout.

REFERENCE

- (1) Rivera, J.: Distribution of Strontium-90 in a 1959 Wheat Sample, *Science*, 133:755-6 (March 17, 1961).

SECTION III.—MILK

Radionuclide Analyses of Pasteurized Milk

PASTEURIZED MILK MONITORING PROGRAM

June 1962

Division of Radiological Health

Division of Environmental Engineering and Food Protection

Public Health Service

Milk monitoring has been conducted by the Public Health Service since early 1957 when the first program was established to develop suitable sampling methods and radiochemical analytical procedures. Milk is produced every day throughout the year in practically every county of the United States. It can be readily checked in processing centers and is a good indicator of fallout in the diet. Raw milk was initially selected for investigation; however, it became evident that the milk actually consumed by the population should be included in a broader sampling program. The result was the initiation, in the first quarter of 1960, of a pasteurized milk sampling program designed to provide data representative of the milk consumed in selected municipalities. Both raw and pasteurized milk sampling data were reported concurrently until June 1961 to permit comparison of the differences between the earlier, limited, milkshed sampling results and those exhibited in the pasteurized milk monitoring.

The June 1961 raw milk sampling results, reported in the November 1961 *Radiological*

Health Data (RHD), represent the last regular publication of such data. A summary discussion of the raw milk sampling program in the December 1961 *RHD* presented the gross relationship between fallout and the occurrence of fission products in milk determined from this study.

During June 1962, the surveillance of pasteurized milk was conducted at 61 stations with the cooperation of State and local milk sanitation agencies who have been shipping samples to the PHS Southwestern, Southeastern, and Northeastern Radiological Health Laboratories for analysis (see figure 1). Publication in *RHD* follows about 3½ months after sample collection because of time required for shipment, processing, decay-product buildup, data compilation, and publication procedures.

The current program emphasizes (1) measurement of the concentrations of radioactivity in samples of pasteurized milk consumed by the public in various regions of the country, and (2) provision of at least one sampling point within virtually all states and additional points




FIGURE 1.—IODINE-131 CONCENTRATION CONTOURS AND MILK SAMPLING STATIONS

when indicated by widely varying conditions of the milk supply or the need to cover large population groups. Each sample is composited in proportion to the volume of milk produced by those plants supplying not less than 90 percent of a city's milk supply. Prior to September 15, 1961, this composite sample was taken from one day's production of pasteurized milk per month and was as representative of a com-

munity's total supply as could be achieved under practical conditions. Since the resumption of nuclear weapons testing, the sampling schedule has been increased in frequency. During June 1962, most stations were sampled twice a week. As all surveillance data are subject to continuing review and evaluation to observe unusual patterns or concentrations which may require immediate attention and







FIGURE 1.—IODINE-131 CONCENTRATION CONTOURS AND MILK SAMPLING STATIONS—Continued

adjustment in the pasteurized milk sampling program operation, further atmospheric nuclear testing may require an immediate re-evaluation and readjustment of the sampling frequency and analytical schedule for this program.

Iodine-131, cesium-137, and barium-140 are determined by gamma scintillation spectroscopy,¹ while strontium-89 and strontium-90 are determined following radiochemical separation. The minimum detectable concentrations in units of $\mu\mu\text{c/liter}$ are: Sr^{89} , 5; Sr^{90} , 1; I^{131} , 10; Cs^{137} , 5; and Ba^{140} , 10.

Table 1 presents summaries of all available analyses for June 1962. When a radionuclide is reported by a laboratory as being below the minimum detectable concentration, one-half of this value is used in calculating the monthly average. A similar procedure is used for the network average.

¹ Southeastern Radiological Health Laboratory employs a radiochemical procedure for barium-140 analysis.

Iodine-131 Geographical Distribution

Just prior to September 1961, the iodine-131 concentrations in pasteurized milk were not detectable. Figure 1 geographically displays the average iodine-131 concentrations in pasteurized milk from September 1961 through June 1962. These plots are based on concentrations in pasteurized milk representing milk sold in 61 cities distributed throughout the United States. Intervening milk sheds for smaller cities or the effect on intra- and interstate milk distribution are not taken into consideration. This permits the use of concentration contours for geographically presenting these data.

During the first few months of 1962, the iodine-131 concentrations in pasteurized milk were not generally detectable, since the iodine-131 present in November and December 1961 had decayed to relatively low levels. In May and June 1962, the concentrations in pasteurized milk again increased due to nuclear

TABLE 1.—RADIOACTIVITY IN PASTEURIZED MILK, JUNE 1962

[Average radioactivity concentrations in $\mu\text{mc/liter}$]

| Sampling location | | Calcium (gm/liter) | | Strontium-90 | | Strontium-90 | | Iodine-131 | | Cesium-137 | | Barium-140 | | Strontium-90 profile in RHD |
|----------------------|----------------|-----------------------|------------------|-------------------|------------------|-------------------|------------------|-------------------|------------------|-------------------|------------------|-------------------|------------------|--------------------------------|
| State | City | Second quarter | Avg for month | Second quarter | Avg for month | Second quarter | Avg for month | Second quarter | Avg for month | Second quarter | Avg for month | Second quarter | Avg for month | |
| Alabama | Montgomery | 1.20 | 1.21 | 105 | 55 | 19 | 16 | 10 | <10 | 55 | 50 | 20 | 20 | Oct. 62 |
| Alaska | Palmer | 1.08 | 1.05 | 35 | 60 | 9 | 13 | 10 | 30 | 20 | 25 | <10 | 10 | |
| Arizona | Phoenix | 1.09 | 1.12 | 15 | 5 | 3 | 3 | <10 | 10 | 10 | 10 | <10 | 10 | Sept. 62 |
| Arkansas | Little Rock | 1.17 | 1.18 | 225 | 120 | 37 | 32 | 20 | 20 | 95 | 80 | 40 | 30 | July 62 |
| California | Sacramento | 1.09 | 1.12 | 30 | 20 | 5 | 5 | 10 | 10 | 15 | 10 | <10 | <10 | Aug. 62 |
| California | San Francisco | 1.10 | 1.07 | 55 | 25 | 9 | 7 | <10 | <10 | 15 | 10 | 10 | <10 | Oct. 62 |
| Colorado | Denver | 1.16 | 1.19 | 30 | 60 | 8 | 13 | <10 | <10 | 10 | 30 | <10 | <10 | Aug. 62 |
| Connecticut | Hartford | 1.19 | 1.24 | 15 | 35 | 11 | 14 | <10 | 10 | 35 | 70 | <10 | <10 | Aug. 62 |
| Delaware | Wilmington | 1.20 | 1.22 | 50 | 40 | 15 | 17 | 10 | 20 | 50 | 80 | <10 | <10 | Sept. 62 |
| District of Columbia | Washington | 1.15 | 1.14 | 50 | 65 | 15 | 17 | <10 | 10 | 40 | 65 | 10 | 20 | Oct. 62 |
| Florida | Tampa | 1.21 | 1.21 | 25 | 25 | 10 | 11 | <10 | 10 | 105 | 120 | <10 | 10 | Sept. 62 |
| Georgia | Atlanta | 1.20 | 1.18 | 120 | 70 | 22 | 20 | <10 | <10 | 80 | 75 | 20 | 20 | Oct. 62 |
| Hawaii | Honolulu | 1.08 | 1.08 | 30 | 25 | 5 | 5 | <10 | 10 | 25 | 30 | <10 | 10 | |
| Idaho | Idaho Falls | 1.09 | 1.12 | 25 | 50 | 7 | 13 | 10 | 30 | 20 | 45 | 10 | 10 | Sept. 62 |
| Illinois | Chicago | 1.16 | 1.19 | 35 | 70 | 11 | 15 | 20 | 30 | 30 | 65 | 10 | 10 | Oct. 62 |
| Indiana | Indianapolis | 1.20 | 1.22 | 50 | 60 | 15 | 16 | 20 | 30 | 40 | 50 | 20 | 20 | Aug. 62 |
| Iowa | Des Moines | 1.08 | 1.09 | 95 | 170 | 15 | 26 | 60 | 70 | 25 | 40 | 20 | 20 | Sept. 62 |
| Kansas | Wichita | 1.08 | 1.08 | 65 | 80 | 12 | 16 | 120 | 130 | 20 | 40 | 20 | 20 | July 62 |
| Kentucky | Louisville | 1.17 | 1.16 | 115 | 120 | 21 | 25 | 20 | 20 | 40 | 60 | 20 | 30 | Sept. 62 |
| Louisiana | New Orleans | 1.23 | 1.24 | 215 | 145 | 37 | 38 | <10 | 10 | 135 | 110 | 30 | 30 | June 62 |
| Maine | Portland | 1.22 | 1.24 | 20 | 45 | 12 | 13 | <10 | 10 | 50 | 95 | <10 | <10 | Sept. 62 |
| Maryland | Baltimore | 1.17 | 1.15 | 45 | 60 | 15 | 17 | 10 | <10 | 50 | 70 | 10 | 20 | Sept. 62 |
| Massachusetts | Boston | 1.16 | 1.13 | 20 | 45 | 13 | 16 | <10 | 10 | 50 | 100 | <10 | <10 | Oct. 62 |
| Michigan | Detroit | 1.17 | 1.18 | 25 | 40 | 11 | 14 | <10 | <10 | 30 | 55 | <10 | <10 | Aug. 62 |
| Michigan | Grand Rapids | 1.23 | 1.21 | 20 | 40 | 10 | 12 | <10 | 10 | 30 | 55 | <10 | 10 | Sept. 62 |
| Minnesota | Minneapolis | 1.08 | 1.10 | 60 | 115 | 13 | 24 | 50 | 30 | 30 | 65 | 10 | 10 | Oct. 62 |
| Mississippi | Jackson | 1.26 | 1.25 | 220 | 140 | 33 | 29 | <10 | 10 | 85 | 85 | 30 | 30 | Aug. 62 |
| Missouri | Kansas City | 1.09 | 1.11 | 130 | 215 | 18 | 25 | 150 | 240 | 30 | 50 | 40 | 50 | Aug. 62 |
| Missouri | St. Louis | 1.08 | 1.12 | 85 | 115 | 16 | 23 | 30 | 60 | 30 | 50 | 10 | 20 | June 62 |
| Montana | Helena | 1.08 | 1.09 | 45 | 75 | 10 | 16 | 20 | 30 | 25 | 50 | 10 | 10 | July 62 |
| Nebraska | Omaha | 1.14 | 1.09 | 70 | 130 | 13 | 24 | 40 | 80 | 20 | 40 | 10 | 20 | Sept. 62 |
| New Hampshire | Manchester | 1.15 | 1.06 | 15 | 35 | 10 | 12 | <10 | <10 | 60 | 115 | <10 | <10 | Sept. 62 |
| New Jersey | Trenton | 1.13 | 1.09 | 25 | 35 | 10 | 12 | <10 | <10 | 30 | 60 | <10 | <10 | Aug. 62 |
| New Mexico | Albuquerque | 1.09 | 1.10 | 15 | 20 | 4 | 6 | 10 | <10 | <5 | <5 | 10 | <10 | July 62 |
| New York | Buffalo | 1.16 | 1.12 | 20 | 40 | 10 | 13 | <10 | 10 | 40 | 80 | <10 | <10 | Aug. 62 |
| New York | New York | 1.13 | 1.14 | 20 | 40 | 12 | 14 | 10 | 10 | 40 | 85 | <10 | <10 | Oct. 62 |
| New York | Syracuse | 1.17 | 1.21 | 20 | 30 | 10 | 12 | 20 | 20 | 25 | 50 | <10 | <10 | July 62 |
| North Carolina | Charlotte | 1.22 | 1.23 | 95 | 60 | 22 | 27 | <10 | 10 | 55 | 70 | 20 | 20 | July 62 |
| North Dakota | Minot | 1.10 | 1.09 | 60 | 120 | 16 | 30 | 20 | 20 | 30 | 70 | <10 | <10 | Sept. 62 |
| Ohio | Cincinnati | 1.18 | 1.21 | 65 | 70 | 17 | 19 | 20 | 20 | 30 | 50 | 20 | 20 | Aug. 62 |
| Ohio | Cleveland | 1.15 | 1.15 | 35 | 80 | 11 | 14 | <10 | 10 | 30 | 65 | <10 | <10 | July 62 |
| Oklahoma | Oklahoma City | 1.15 | 1.14 | 115 | 135 | 21 | 29 | 60 | 100 | 60 | 95 | 30 | 50 | Oct. 62 |
| Oregon | Portland | 1.13 | 1.14 | 95 | 65 | 17 | 14 | 10 | 30 | 55 | 65 | 10 | <10 | Aug. 62 |
| Pennsylvania | Philadelphia | 1.16 | 1.17 | 35 | 40 | 13 | 13 | <10 | 10 | 45 | 75 | <10 | <10 | July 62 |
| Pennsylvania | Pittsburgh | 1.17 | 1.18 | 35 | 55 | 14 | 17 | 10 | 30 | 45 | 90 | <10 | <10 | June 62 |
| Puerto Rico | San Juan | 1.13 | 1.15 | 75 | 55 | 11 | 13 | 20 | 20 | 55 | 50 | 20 | 20 | |
| Rhode Island | Providence | 1.19 | 1.23 | 20 | 35 | 12 | 13 | 10 | 10 | 45 | 85 | <10 | <10 | Sept. 62 |
| South Carolina | Charleston | 1.20 | 1.19 | 95 | 60 | 21 | 20 | <10 | <10 | 75 | 80 | 20 | 20 | Aug. 62 |
| South Dakota | Rapid City | 1.09 | 1.19 | 70 | 185 | 17 | 40 | <10 | <10 | 30 | 85 | 10 | <10 | Oct. 62 |
| Tennessee | Chattanooga | 1.24 | 1.22 | 200 | 105 | 28 | 24 | 10 | <10 | 90 | 70 | 30 | 20 | Oct. 62 |
| Tennessee | Memphis | 1.21 | 1.20 | 160 | 115 | 27 | 26 | <10 | 10 | 40 | 45 | 30 | 20 | Sept. 62 |
| Texas | Austin | 1.17 | 1.14 | 35 | 30 | 8 | 8 | <10 | <10 | 25 | 40 | <10 | 10 | Oct. 62 |
| Texas | Dallas | 1.18 | 1.19 | 115 | 85 | 18 | 18 | 20 | 30 | 50 | 50 | 20 | 30 | Aug. 62 |
| Utah | Salt Lake City | 1.11 | 1.12 | 25 | 40 | 8 | 14 | <10 | 10 | 30 | 75 | <10 | 10 | June 62 |
| Vermont | Burlington | 1.13 | 1.10 | 15 | 30 | 10 | 13 | <10 | <10 | 30 | 55 | <10 | <10 | Aug. 62 |
| Virginia | Norfolk | 1.19 | 1.19 | 70 | 70 | 17 | 21 | <10 | 10 | 55 | 80 | 10 | 20 | Aug. 62 |
| Washington | Seattle | 1.12 | 1.15 | 80 | 65 | 18 | 19 | <10 | 10 | 55 | 75 | <10 | <10 | Oct. 62 |
| Washington | Spokane | 1.13 | 1.16 | 55 | 85 | 14 | 18 | 120 | 350 | 35 | 55 | <10 | <10 | Aug. 62 |
| West Virginia | Charleston | 1.17 | 1.14 | 80 | 115 | 18 | 23 | 20 | 20 | 35 | 60 | 20 | 30 | Oct. 62 |
| Wisconsin | Milwaukee | 1.20 | 1.18 | 15 | 35 | 8 | 11 | 10 | 20 | 15 | 25 | <10 | <10 | Sept. 62 |
| Wyoming | Laramie | 1.09 | 1.13 | 35 | 60 | 8 | 15 | 20 | 20 | 25 | 50 | 10 | 10 | Sept. 62 |
| Network average | | 1.15 | 1.16 | 65 | 71 | 14 | 17 | 20 | 30 | 40 | 61 | 10 | 14 | |

weapons testing. The highest June average iodine-131 concentration in pasteurized milk appeared at Spokane, Washington.

Strontium-90 Profiles

In previous issues of *RHD*, the average monthly strontium-90 concentrations in pas-

teurized milk from selected cities in the monitoring program were published. An additional 14 cities are presented in figure 2. Each graph shows the strontium-90 concentrations in milk from selected cities in 4 geographic regions of the United States. The usual




FIGURE 2.—STRONTIUM-90 CONCENTRATIONS IN PASTEURIZED MILK

regional differences (higher strontium-90 in the East than in the West) are not evident in the selected profiles. The North Central region

has shown the greatest change. Strontium-90 concentrations in milk produced in this region are now similar to those prevalent in the South.

Radionuclides in Raw Milk from the Vicinity in the Nevada Test Site

July 1961-June 1962

U.S. Atomic Energy Commission

Shortly after the Operation Plumbob test series in 1957, the Atomic Energy Commission Off-Site Organization at the Nevada Test Site contracted with the Public Health Service's Robert A. Taft Sanitary Engineering Center for analyses of radionuclides in raw milk (Las Vegas milk is pasteurized) produced in the vicinity of the Nevada Test Site. In April 1960, the Southwestern Radiological Health Laboratory of the Public Health Service Division of Radiological Health assumed responsibility for the radionuclide analyses of milk samples obtained in the off-site monitoring program (see sampling locations on figure 1). All milk sample analyses taken in the vicinity of the



FIGURE 1.—MILK SAMPLING LOCATIONS AT THE NEVADA TEST SITE

TABLE 1.—RADIONUCLIDE CONCENTRATIONS IN MILK SAMPLES FROM THE VICINITY OF NEVADA TEST SITE

[Radioactivity concentrations in $\mu\text{c}/\text{liter}$]

| Collection date | Location | Calcium (g/liter) | Strontium-89 | Strontium-90 | Iodine-131 | Cesium-137 | Barium-140 lanthanum-140 |
|-----------------|--------------------------|----------------------|--------------|--------------|------------|------------|-----------------------------|
| July 20, 1961 | Las Vegas | 0.94 | <5 | 2 | <10 | <10 | <20 |
| Aug. 21, 1961 | Las Vegas | 0.86 | <5 | 2 | <10 | <10 | <20 |
| Aug. 15, 1961 | Alamo | 0.86 | <5 | 2 | <10 | <10 | <20 |
| Sept. 6, 1961 | Springdale | a | a | a | <10 | <10 | 570 |
| Oct. 31, 1961 | Hiko | 1.08 | 95 | 4 | 720 | 20 | 60 |
| Oct. 31, 1961 | Las Vegas | 1.10 | 15 | 3 | 40 | <10 | <20 |
| Nov. 1, 1961 | Glendale | 1.06 | <5 | 2 | 60 | <10 | <20 |
| Nov. 2, 1961 | Caliente | 1.20 | <5 | 4 | 80 | <10 | 20 |
| Nov. 29, 1961 | Fallini's Ranch | 1.33 | 40 | 6 | 90 | 50 | 70 |
| Dec. 1, 1961 | Las Vegas | 1.02 | 15 | 2 | 40 | 25 | 60 |
| Dec. 1, 1961 | Diabla | 1.16 | 5 | 3 | 30 | 15 | <20 |
| Dec. 13, 1961 | Alamo | 1.07 | 10 | 2 | <10 | <10 | <20 |
| Dec. 13, 1961 | Hiko | 1.10 | <5 | 2 | <10 | <10 | <20 |
| Dec. 18, 1961 | Las Vegas | 1.07 | <5 | 4 | <10 | <10 | <20 |
| Dec. 21, 1961 | Las Vegas | 1.12 | 5 | 3 | <10 | <10 | <20 |
| Jan. 3, 1962 | Panaca | 1.10 | <5 | 2 | <10 | <10 | <20 |
| Feb. 13, 1962 | Las Vegas | 1.00 | <5 | 2 | <10 | 10 | <20 |
| Feb. 22, 1962 | Alamo | 1.00 | <5 | 2 | <10 | <10 | <20 |
| Feb. 26, 1962 | Springdale | 1.00 | 30 | 3 | <10 | 30 | <20 |
| Apr. 4, 1962 | Caliente | 1.10 | 5 | 3 | <10 | <10 | <20 |
| Apr. 18, 1962 | Fallini's Ranch | 1.18 | 10 | 5 | <10 | 10 | <20 |
| Apr. 19, 1962 | Alamo | 1.15 | <5 | 3 | <10 | <10 | <20 |
| Apr. 19, 1962 | Panaca | 1.14 | 5 | 4 | <10 | 10 | <20 |
| Apr. 25, 1962 | Lund | 1.08 | <5 | 4 | <10 | <10 | <20 |
| Apr. 26, 1962 | Springdale | 1.08 | 25 | 4 | 20 | 25 | <20 |
| May 9, 1962 | Lund | 1.08 | <5 | 1 | <10 | <10 | <20 |
| May 17, 1962 | Hiko | 1.14 | 5 | 1 | <10 | <10 | <20 |
| June 5, 1962 | Searles Ranch | b | b | b | <10 | 20 | <20 |
| June 5, 1962 | Las Vegas | b | b | b | <10 | 30 | <20 |
| June 5, 1962 | Caliente | b | b | b | <10 | 25 | <20 |
| June 5, 1962 | Alamo | b | b | b | <10 | 35 | <20 |
| June 7, 1962 | Springdale | b | b | b | <10 | 50 | <20 |
| June 4, 1962 | Panaca | b | b | b | <10 | <10 | <20 |
| June 8, 1962 | Fallini's Ranch | b | b | b | <10 | 70 | <20 |
| June 13, 1962 | White River Valley Ranch | b | b | b | 40 | 80 | <20 |
| June 18, 1962 | Blue Eagle Ranch | b | b | b | 20 | 180 | <20 |
| June 20, 1962 | Adavan | b | b | b | 360 | 75 | <20 |
| June 21, 1962 | Elko | b | b | b | 610 | 200 | <20 |
| June 22, 1962 | Robbins Ranch | b | b | b | 520 | 140 | <20 |
| June 22, 1962 | Carlin | b | b | b | 160 | 120 | <20 |
| June 23, 1962 | Wells | b | b | b | 80 | 110 | <20 |

a Sample not valid.

b Not available in time for inclusion.

Nevada Test Site since July 20, 1961, are reported in table 1. Gross beta concentrations determined by this monitoring program in

water supplies in off-site areas of the Nevada Test Site have been published previously in *Radiological Health Data*.

Strontium-90 in Canadian Dried Milk Products

RADIOACTIVE FALLOUT STUDY PROGRAM

First Quarter 1962

Department of National Health and Welfare, Ottawa, Canada

Beginning November 1955, radiochemical analyses of skim milk and buttermilk powders for strontium-90 concentrations were initiated by the Department of National Health and Welfare. Through the cooperation of the Marketing Division of the Canadian Department of Agriculture, four 1-pound samples of dried milk from each station were collected by dairy products inspectors each month for the Radioactive Fallout Study Program. No precise control is maintained over the method of sample collection, since whether or not all samples are collected on the same day or if the sample is truly representative of the milk produced in a given area is not known.

Because of uncertainties introduced by the method of sampling, the significance of differences between station-to-station and month-to-

month results is not precisely known. It is possible, however, to consider all the results for a given period of time as being sufficiently random in selection to show any national trend when average values for such periods and all stations are plotted, as in figure 2. It should be emphasized that average concentrations over a considerable period of time are more meaningful from the public health point of view than fluctuating monthly concentrations.

A detailed discussion of the sampling and radiochemical methodology employed may be found in the Department's publication (1).

Table 1 presents the results of measurements of strontium-90 in Canadian dried milk powder for the first quarter of 1962, which were taken from the "Quarterly Report of the Radioactive Fallout Monitoring Program," dated July 1961,

TABLE 1.—STRONTIUM-90 IN CANADIAN DRIED MILK POWDER, 1962

[Concentrations in $\mu\text{c/g Ca}$]

| Station | Type of milk | January | February | March |
|-----------------------------------------------|--------------|---------|----------|-------|
| Calgary..... | Skim milk | 4.8 | 4.6 | 5.0 |
| Edmonton..... | Skim milk | 6.6 | 7.3 | 5.3 |
| Granby region: | | | | |
| Lawrenceville..... | Skim milk | 14.0 | 15.4 | 14.9 |
| Nicolet..... | Skim milk | 6.0 | 7.5 | 8.3 |
| Halifax..... | Skim milk | 7.7 | 7.0 | 9.0 |
| La Durantye..... | Skim milk | 11.5 | 11.2 | N.S. |
| London..... | Skim milk | 3.4 | 3.3 | 4.6 |
| Megantic..... | Skim milk | 13.0 | 10.3 | 8.8 |
| Moncton..... | Buttermilk | 7.5 | 6.7 | 7.7 |
| Ottawa..... | Skim milk | 4.7 | 5.1 | 6.1 |
| Saskatoon..... | Buttermilk | 4.7 | 4.5 | 6.9 |
| Sussex..... | Skim milk | 8.2 | 9.4 | 9.1 |
| Vancouver..... | Skim milk | 7.7 | 7.0 | 6.7 |
| Walkerton..... | Skim milk | 4.6 | 4.4 | 3.2 |
| Winnipeg region: | | | | |
| Arborg..... | Buttermilk | 4.8 | 4.8 | 4.3 |
| Grunthal..... | Buttermilk | 4.6 | 4.5 | 4.0 |
| Average..... | | 7.1 | 7.1 | 6.9 |
| 12-month average (April 1961–March 1962)..... | | | | 0.8 |



FIGURE 1.—CANADIAN DRIED MILK SAMPLING STATIONS

published by the Radiation Protection Division of the Department of National Health and Welfare.

For an approximate method to compare the Canadian powdered milk data in units of $\mu\text{C/g Ca}$ to other milk data in units of $\mu\text{C/liter}$, it can be assumed that Canadian powdered milk has approximately 1.2 g Ca/liter of liquid milk. Thus, Canadian powdered milk data may be compared to liquid milk by multiplying the former values by 0.8.




FIGURE 2.—AVERAGE STRONTIUM-90 CONCENTRATIONS IN CANADIAN DRIED MILK PRODUCTS

Previous coverage in *Radiological Health Data*:

| Period | Issue |
|-------------------------|----------------|
| January 1956–March 1960 | September 1960 |
| Second quarter 1960 | December 1960 |
| Third quarter 1960 | April 1961 |
| Fourth quarter 1960 | August 1961 |
| First quarter 1961 | December 1961 |
| Second quarter 1961 | February 1962 |
| Third quarter 1961 | May 1962 |
| Fourth quarter 1961 | September 1962 |

REFERENCE

- (1) Radiation Protection Division, Department of National Health and Welfare, Ottawa, Canada: Preliminary Report on the Measurements of Radioactive Strontium in Canadian Milk Powder Samples, CNHW (RP-1), (July 1958).

Iodine-131 and Strontium-90 in Milk from the United Kingdom^{1, 2}

January 1961–April 1962

Agricultural Research Council Radiobiological Laboratory, England

Since 1958, the Agricultural Research Council Radiobiological Laboratory has performed strontium-90 analyses of raw milk produced in the United Kingdom. For the "country-wide" survey, raw milk samples are obtained every two weeks from over 200 depots selected according to a statistical plan, which handle more than 40 percent of the country's total milk production. The milk samples from the depots are composited to represent each zone as shown in figure 1.

¹ Agricultural Research Council Radiobiological Laboratory: *Interim Report on Radioactivity in Milk*, ARCRL-6, Her Majesty's Stationery Office, London (December 1961), price 2 s.Od.

² Agricultural Research Council Radiobiological Laboratory: *Interim Report on Radioactivity in Diet*, ARCRL-7, Her Majesty's Stationery Office, London (May 1962), price 2 s.Od.

Iodine-131

When it became apparent in September 1961 that the resumption of nuclear weapons testing was likely to lead to an appreciable contamination of milk, arrangements were made to receive milk daily because considerable day to day variations can occur in the iodine-131 concentrations in milk. Thus, during the period when the highest levels occurred (October and November, 1961) samples were obtained daily from up to 23 "regular sampling points" of which at least two were in each of the zones listed in table 1. In addition, country-wide sampling was also carried out once each week to provide fuller information on regional differences. When the levels decreased in early December 1961, the number of regular sampling



FIGURE 1.—MILK SAMPLING LOCATIONS FOR COUNTRY-WIDE SURVEY

points was reduced to 7 and later to 6, distributed throughout the country. The frequency of the collections from the country-wide survey was reduced to once every 2 weeks. By the end of December the amount of iodine-131 in milk was below the limit of accurate detection in all samples, therefore the number of analyses was further reduced by bulking milk collected from wider areas.

The iodine-131 concentrations were determined by gamma counting after a rapid concentration process. Formalin is added to the milk and the proteins precipitated by the addition of trichloroacetic acid; silver halide is subsequently precipitated. The gamma activity of the combined precipitate of protein and silver chloride is then counted using a 4" by 3" sodium iodide (thallium activated) crystal; the content of iodine-131 is calculated from the count-rate in the 364 kev photopeak.

The iodine-131 results from the country-wide survey and the regular sampling points have been combined in table 1. Because of the magnitude of day to day variations (see figure 2) the results of the country-wide survey do not by themselves provide a precise estimate of the




FIGURE 2.—IODINE-131 IN UNITED KINGDOM MILK

average level since sampling was commenced. Some samples were collected on days when the results for the regular sampling points show that the content of milk was either appreciably higher or lower than the weekly average. A better estimate of the weekly mean levels throughout the country can be obtained by combining the results for the regular sampling sites with those for the country-wide survey. This was done by applying a correction factor to the country-wide samples consisting of the ratio of the mean value for the regular sampling points on the same day the country-wide sample was taken to the mean value for those points during each week. This calculation was made separately for each zone. For the 10 zones listed in table 1 the mean concentration for the 32 weeks ending April 28, 1962, ranged from 23 to 51 pc/liter, with a country-wide average level of 36 pc/liter.

Strontium-90

Whereas iodine-131 can be measured in milk within a few hours of the receipt of samples, measurements of strontium-90 take a considerably longer period. Not only is elaborate chemical separation necessary but also the strontium-90 can be adequately determined only by studying the decay of its radioactive

daughter, yttrium-90, over a period of several weeks. Therefore, the reporting of strontium-90 analyses lags that for iodine-131 by at least 2 months. January through December 1961 strontium-90 analyses are reported in table 2.

During 1961 the collection of milk for the country-wide survey of strontium-90 contamination was continued on the basis described previously. On account of the marked reduction in the concentrations of strontium-90 in

milk during 1960 it was considered justified to reduce the analytical effort given to this work. The milk from 5 zones in England which had previously been assayed separately every 3 months was combined for analysis; after weapons testing was resumed in September 1961, the milk from each zone was again examined separately. In May 1961, the temporary division of Scotland into two zones was abandoned.

TABLE 1.—IODINE-131 CONCENTRATIONS IN MILK THROUGHOUT THE UNITED KINGDOM, 1961-1962

[Average concentrations in pc/liter]

| Zone (See figure 1) | September | | October | | | | November | | | | December | | | | Apr | Average Sept-Apr |
|------------------------------|-----------|-------|---------|------|-------|-------|----------|------|-------|-------|----------|-----|-------|-------|-------|---------------------|
| | 16-23 | 24-30 | 1-7 | 8-14 | 15-21 | 22-28 | 29-4 | 5-11 | 12-18 | 19-25 | 26-2 | 3-9 | 10-16 | 17-24 | 25-28 | 16-28 |
| | | | | | | | | | | | | | | | | |
| 1. Northern England | 76 | 97 | 123 | 115 | 87 | 117 | 119 | 88 | 100 | 75 | 27 | <15 | <15 | <15 | <15 | 32 |
| 2. Northwest England | 57 | 73 | 93 | 131 | 99 | 109 | 115 | 102 | 97 | 73 | 43 | 18 | <15 | <15 | <15 | 32 |
| 3. East & East Midlands | 52 | 66 | 84 | 78 | 59 | 100 | 81 | 56 | 59 | 66 | 44 | <15 | <15 | <15 | <15 | 23 |
| 4. West Midlands | 67 | 86 | 108 | 130 | 98 | 111 | 119 | 121 | 118 | 76 | 31 | 31 | 41 | 23 | <15 | 36 |
| 5. Wales | 89 | 114 | 145 | 234 | 177 | 201 | 194 | 159 | 155 | 87 | 50 | 24 | <15 | <15 | <15 | 51 |
| 6. South & Southeast England | 83 | 107 | 136 | 88 | 67 | 84 | 125 | 57 | 84 | 65 | 50 | 24 | <15 | <15 | <15 | 30 |
| 7. Midwestern England | 56 | 72 | 91 | 114 | 86 | 108 | 141 | 124 | 102 | 89 | 50 | 21 | <15 | <15 | <15 | 33 |
| 8. Farwestern England | 52 | 67 | 85 | 187 | 142 | 205 | 175 | 143 | 138 | 79 | 43 | 37 | 37 | 21 | <15 | 44 |
| 9. Scotland | 139 | 179 | 227 | 208 | 157 | 143 | 104 | 93 | 84 | 30 | 18 | 18 | <15 | <15 | <15 | 44 |
| 10. Northern Ireland | 132 | 169 | 214 | 152 | 115 | 156 | 163 | 200 | 129 | 77 | 32 | 21 | 34 | 19 | <15 | 50 |
| Production weighted mean | 77 | 99 | 125 | 141 | 106 | 127 | 130 | 108 | 103 | 71 | 40 | 20 | <15 | <15 | <15 | 36 |

TABLE 2.—STRONTIUM-90 CONCENTRATIONS IN MILK THROUGHOUT THE UNITED KINGDOM, 1961

[Average concentrations in pc/liter]

| Zone (See figure 1) | January 1- February 1 | February 15- March 15 | April 1- May 1 | May 15- June 15 | July 1- August 1 | August 15- September 15 | October 1- November 1 | November 15- December 15 |
|------------------------------------------------|--------------------------|--------------------------|-------------------|--------------------|---------------------|----------------------------|--------------------------|-----------------------------|
| 1. Northern England..... | a— | — | — | — | — | — | b6.8 | b6.8 |
| 2. Northwest England..... | 6.7 | 5.5 | 6.7 | 6.4 | 5.8 | 5.0 | 5.2 | 6.4 |
| 3. East & East Midlands..... | — | — | — | — | — | — | b3.7 | b3.7 |
| 4. West Midlands..... | — | — | — | — | — | — | b5.2 | b5.2 |
| 5. Wales..... | 9.4 | 9.0 | 10.4 | 10.01 | 8.4 | 8.2 | 10.4 | 10.5 |
| 6. South & Southeast England..... | — | — | — | — | — | — | b4.2 | b4.2 |
| 7. Midwestern England..... | 6.6 | 5.1 | 5.7 | 6.0 | 4.6 | 5.7 | 5.5 | 5.6 |
| 8. Far Western England..... | — | — | — | — | — | — | b8.8 | b8.8 |
| 9.1 Southwest Scotland..... | 8.4 | 8.6 | 9.2 | 9.2 | 9.8 | 8.4 | 10.0 | 7.4 |
| 9.2 North & East Scotland..... | 7.4 | 7.4 | 6.9 | — | — | — | — | — |
| 10. Northern Ireland..... | 11.0 | 10.8 | 11.4 | 10.8 | 9.0 | 9.6 | 9.6 | 11.6 |
| Remainder of England..... | b6.3 | b6.3 | b6.4 | b6.4 | b6.2 | b6.2 | — | — |
| Production weighted mean for latest 12 months. | | 6.93 | | 6.58 | | 6.65 | | 6.82 |

^a Dash indicates sample composited with 4 other sections. Analysis is reported as "Remainder of England."

^b Analysis covers 2 time periods.

SECTION IV.—WATER

Radioactivity in Raw Surface Waters

NATIONAL WATER QUALITY NETWORK April 1962

Division of Water Supply and Pollution Control, Public Health Service

The National Water Quality Network, operated in cooperation with State and local agencies, commenced operations in October 1957 (1). By the end of April 1962, 108 stations located on major waterways used for public water supply, propagation of fish and wildlife, and recreational, industrial, and agricultural purposes were submitting water samples for analyses of plankton population, organic chemicals, stream flow data, chemical, biological, and physical quality, and radioactivity (2, 3). Samples are taken weekly, monthly, or continually, depending on the type of analysis and the degree of water quality.

The radioactivity associated with dissolved solids provides a rough measure of the levels which may be found in treated water, since nearly all of the suspended matter is removed by treatment procedures (4). Naturally-occurring radioactive substances in the environment are the source of essentially all the alpha activity, while contamination from man-made sources is the major contributor of beta activity. Natural beta activity can be two or three times the natural alpha activity based on the presence of the same nuclides. Gross alpha and beta measurements are made on both suspended

and dissolved solids (strontium-90 on the total solids only) in raw surface water samples according to standard procedures (5, 6).

For the first two years, alpha determinations were reported generally on composites of more than one weekly sample. As of January 1960, the policy was established that analyses were to be done weekly for the first operating year of each new station. At some stations on the Colorado and Animas Rivers, determinations were done on weekly samples or semimonthly on 2- or 3-week composites. The remainder of the stations were scheduled so that each station made one gross alpha determination per month. Beginning with November 1961, gross alpha determinations have been made on one sample from each station each month, unless there is evidence of alpha activity in excess of background levels. In the latter instance, an alpha determination has been made on a weekly or biweekly basis, depending on the particular station.

For the first two years of the network operations, beta determinations were made on weekly samples. Since January 1959, a portion of each sample from all stations in the network has been composited into a 3-month station sample

TABLE 1.—RADIOACTIVITY IN RAW SURFACE WATERS

[Average concentrations in $\mu\text{c}/\text{liter}$]

| Station | Quarter ending Dec. 31, 1961 | April 1962 | | | | | | |
|-----------------------------------------|---------------------------------------|------------------|---------------|-----------|-----------|----------------|-----------|-----------|
| | | Strontium- 90 | Beta Activity | | | Alpha activity | | |
| | | | Total | Suspended | Dissolved | Total | Suspended | Dissolved |
| Allegheny River: Pittsburgh, Pa. | — | 13 | 8 | 21 | — | — | — | — |
| Animas River: Cedar Hill, N. Mex. | 0.9 | 71 | 30 | 101 | 17 | 1 | 18 | 18 |
| Apalachicola River: Chattahoochee, Fla. | — | 30 | 24 | 54 | 1 | 0 | 1 | 1 |
| Arkansas River: | | | | | | | | |
| Coolidge, Kansas | 1.0 | 19 | 62 | 81 | 2 | 15 | 17 | 17 |
| Ponca City, Okla. | — | 21 | 54 | 75 | — | — | — | — |
| Pendleton Ferry, Ark. | — | 87 | 37 | 124 | 2 | 2 | 4 | 4 |
| Big Horn River: Hardin, Mont. | — | 47 | 34 | 81 | 9 | 4 | 13 | 13 |
| Big Sioux River: Sioux Falls, So. Dak. | 1.2 | 26 | 74 | 100 | 2 | 2 | 4 | 4 |
| Chattahoochee River: Atlanta, Ga. | 0.3 | 15 | 8 | 23 | 0 | 0 | 0 | 0 |
| Clear Water River: East Lewiston, Idaho | 0.3 | 19 | 10 | 29 | 1 | 0 | 1 | 1 |
| Colorado River: | | | | | | | | |
| Loma, Colo. | — | 86 | 51 | 137 | 10 | 6 | 16 | 16 |
| Boulder City, Nev. | 1.8 | 5 | 18 | 23 | 0 | 7 | 7 | 7 |
| Parker Dam, Calif. | — | 0 | 13 | 13 | 0 | 5 | 5 | 5 |
| Yuma, Ariz. | — | 17 | 32 | 49 | — | — | — | — |
| Columbia River: | | | | | | | | |
| Wenatchee, Wash. | 0.7 | 2 | 9 | 11 | — | — | — | — |
| Pasco, Wash. | — | 72 | 237 | 309 | — | — | — | — |
| McNary Dam, Oreg. | — | 50 | 210 | 260 | 0 | 1 | 1 | 1 |
| Connecticut River: | | | | | | | | |
| Northfield, Mass. | — | 29 | 21 | 50 | — | — | — | — |
| Wilder, Vt. | 3.7 | 12 | 20 | 32 | 0 | 0 | 0 | 0 |
| Cumberland River: Clarksville, Tenn. | 0.9 | — | — | — | — | — | — | — |
| Delaware River: | | | | | | | | |
| Philadelphia, Pa. | — | 23 | 24 | 47 | — | — | — | — |
| Trenton, New Jersey | 0.7 | 10 | 19 | 29 | 0 | 0 | 0 | 0 |
| Martins Creek, Pa. | — | 13 | 16 | 29 | 0 | 0 | 0 | 0 |
| Ecumbia River: Century, Fla. | — | 39 | 27 | 66 | — | — | — | — |
| Great Lakes: | | | | | | | | |
| Buffalo, N. Y. | 0.8 | 12 | 16 | 28 | 0 | 0 | 0 | 0 |
| Detroit, Mich. | — | 3 | 11 | 14 | 0 | 0 | 0 | 0 |
| Port Huron, Mich. | — | 7 | 7 | 14 | — | — | — | — |
| Sault Ste. Marie, Mich. | — | 2 | 3 | 5 | — | — | — | — |
| Gary, Ind. | 0.1 | 10 | 6 | 16 | 0 | 0 | 0 | 0 |
| Duluth, Minn. | 0.1 | 2 | 4 | 6 | 0 | 0 | 0 | 0 |
| Hudson River: Poughkeepsie, N. Y. | 2.5 | — | — | — | — | — | — | — |
| Illinois River: | | | | | | | | |
| Grafton, Ill. | — | 110 | 57 | 167 | 13 | 0 | 13 | 13 |
| Peoria, Ill. | — | 28 | 39 | 67 | — | — | — | — |
| Kanawha River: Winfield Dam, W. Va. | — | 7 | 15 | 22 | — | — | — | — |
| Klamath River: Copco, Oreg. | 1.2 | 18 | 28 | 46 | — | — | — | — |
| Little Miami River: Cincinnati, Ohio | 0.6 | 7 | 37 | 44 | 1 | 0 | 1 | 1 |
| Mississippi River: | | | | | | | | |
| St. Paul, Minn. | — | 40 | 66 | 106 | 0 | 1 | 1 | 1 |
| Dubuque, Iowa | — | 44 | 22 | 66 | 2 | 0 | 2 | 2 |
| Burlington, Iowa | — | 58 | 37 | 95 | 9 | 0 | 9 | 9 |
| E. St. Louis, Ill. | — | 48 | 42 | 90 | 3 | 0 | 3 | 3 |
| Cape Girardeau, Mo. | — | 66 | 40 | 106 | 12 | 2 | 14 | 14 |
| Delta, La. | — | 69 | 38 | 107 | 3 | 1 | 4 | 4 |
| New Orleans, La. | — | 78 | 26 | 104 | 4 | 1 | 5 | 5 |
| Vicksburg, Miss. | — | 56 | 31 | 87 | 2 | <1 | 2 | 2 |
| Missouri River: | | | | | | | | |
| Williston, N. Dak. | — | 38 | 19 | 57 | — | — | — | — |
| Bismarck, N. Dak. | — | 14 | 25 | 39 | 1 | 4 | 5 | 5 |
| Yankton, S. Dak. | 2.5 | 64 | 82 | 146 | — | — | — | — |
| Omaha, Nebr. | — | 38 | 63 | 101 | — | — | — | — |
| St. Joseph, Mo. | — | 254 | 49 | 303 | — | — | — | — |
| Kansas City, Kans. | — | 277 | 57 | 334 | — | — | — | — |
| Missouri City, Mo. | — | 164 | 39 | 203 | 19 | 2 | 21 | 21 |
| St. Louis, Mo. | 2.1 | 124 | 48 | 172 | — | — | — | — |
| Monongahela River: Pittsburgh, Pa. | — | 4 | 13 | 17 | <1 | <1 | <1 | <1 |
| North Platte River: Henry, Nebr. | 1.3 | 17 | 50 | 67 | 1 | 31 | 32 | 32 |
| Ohio River: | | | | | | | | |
| East Liverppol, Ohio | — | 11 | 25 | 36 | 2 | 0 | 2 | 2 |
| Huntington, W. Va. | — | 32 | 19 | 51 | — | — | — | — |
| Cincinnati, Ohio | — | 55 | 17 | 72 | 3 | 0 | 3 | 3 |
| Louisville, Ky. | — | 46 | 16 | 62 | 4 | 1 | 5 | 5 |
| Evansville, Ind. | — | 46 | 15 | 61 | — | — | — | — |
| Cairo, Ill. | — | 158 | 14 | 172 | 14 | 0 | 14 | 14 |
| Ouachita River: Bastrop, La. | 0.7 | 40 | 52 | 92 | 0 | <1 | <1 | <1 |
| Platte River: Plattsmouth, Nebr. | — | 83 | 54 | 137 | 16 | 6 | 22 | 22 |
| Potomac River: | | | | | | | | |
| Williamsport, Md. | 0.9 | 12 | 14 | 26 | 0 | 0 | 0 | 0 |
| Great Falls, Md. | — | 18 | 16 | 34 | — | — | — | — |
| Rainy River: | | | | | | | | |
| Baudette, Minn. | — | 2 | 14 | 16 | 1 | 0 | 1 | 1 |
| International Fls, Minn. | — | 2 | 12 | 14 | 0 | 0 | 0 | 0 |
| Red River, South: | | | | | | | | |
| Alexandria, La. | — | 124 | 46 | 170 | 6 | 0 | 6 | 6 |
| Denison, Tex. | — | 29 | 27 | 56 | 1 | 1 | 2 | 2 |
| Rio Grande River: | | | | | | | | |
| Alamosa, Colo. | — | 29 | 26 | 55 | 1 | 2 | 3 | 3 |
| El Paso, Tex. | — | 39 | 22 | 61 | — | — | — | — |
| Laredo, Tex. | — | 13 | 45 | 58 | 1 | 3 | 4 | 4 |
| Brownsville, Tex. | — | 3 | 27 | 30 | — | — | — | — |

TABLE 1.—RADIOACTIVITY IN RAW SURFACE WATERS—Continued

[Average concentrations in $\mu\text{mc/liter}$]

| Station | Quarter ending Dec. 31, 1961 | April 1962 | | | | | |
|----------------------------------------------|------------------------------|--------------|---------------|-----------|-------|----------------|-----------|
| | | Strontium-90 | Beta Activity | | | Alpha activity | |
| | | | Suspended | Dissolved | Total | Suspended | Dissolved |
| | | Total | | | | | Total |
| Roanoke River: John H. Kerr Resr. & Dam, Va. | — | — | 36 | 20 | 56 | <1 | 0 |
| Sabine River: Ruliff, Tex. | — | — | 23 | 31 | 54 | — | — |
| San Juan River: Shiprock, N. Mex. | — | — | 123 | 26 | 149 | 14 | 2 |
| St. Lawrence River: Massena, N. Y. | 1.6 | — | 16 | 20 | 36 | 1 | 0 |
| Schuylkill River: Philadelphia, Penn. | — | — | 9 | 15 | 24 | — | — |
| Savannah River: Port Wentworth, Ga. | — | — | 16 | 27 | 43 | 0 | 0 |
| Shenandoah River: Berryville, Va. | 0.2 | — | 30 | 13 | 43 | 1 | 0 |
| Snake River: | | | | | | | |
| Wawawai, Wash. | — | — | 12 | 14 | 26 | — | — |
| Payette, Idaho. | — | — | 30 | 26 | 56 | 1 | 2 |
| South Platte River: Julesburg, Colo. | — | — | 40 | 46 | 86 | 3 | 19 |
| Susquehanna River: | | | | | | | |
| Sayre, Pa. | — | — | 30 | 15 | 45 | — | — |
| Conowingo, Md. | — | — | 26 | 16 | 42 | 3 | 0 |
| Tennessee River: | | | | | | | |
| Lenoir City, Tenn. | — | — | 13 | 14 | 27 | 0 | 0 |
| Chattanooga, Tenn. | 0.5 | — | 24 | 58 | 82 | 1 | 0 |
| Pickwick Landing, Tenn. | — | — | 31 | 38 | 69 | 0 | 0 |
| Tombigbee River: Columbus, Miss. | 0.2 | — | — | — | — | — | — |
| Truckee River: Farad, Calif. | — | — | 28 | 15 | 43 | 1 | 0 |
| Verdigris River: Nowata, Okla. | — | — | 20 | 40 | 60 | 0 | 2 |
| Yakima River: Richland, Wash. | — | — | 16 | 11 | 27 | <1 | <1 |
| Yellowstone River: Sidney, Mont. | — | — | 46 | 33 | 79 | — | — |

for measurement of strontium-90 (7). Beginning January 1960, analyses were performed weekly for the first operating year of each new station. Weekly analyses were to be continued indefinitely for all stations which may be affected by waste discharges from nuclear installations. Where stations have been operated for more than 1 year, semimonthly determinations on composites of 2 or 3 weekly samples were performed when beta activity was found to be above background. Monthly determinations on composites of all samples received from a station during the month were conducted on samples from streams where beta activity was at background levels. Since September 1961, gross beta determinations have been made on all samples collected.

The data reported in table 1 represent the average of all data reported for the periods indicated. The reported strontium-90 data are the results of determinations on 3-month composite samples for the quarter ending in the month shown.

REFERENCES

- (1) Public Health Service: National Water Quality Control Network, *Fallout From Nuclear Weapons Tests*, 1:167-9, Hearings before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, Superintendent of Documents, U.S. Government Printing Office, Washington 25, D.C. (May 1959).
- (2) Division of Water Supply and Pollution Control, Public Health Service: "National Water Quality Network Annual Compilation of Data," PHS publication No. 663, 1960 Edition, Superintendent of Documents, U.S. Government Printing Office, Washington 25, D.C.
- (3) Setter, L. R., and S. L. Baker: Radioactivity of Surface Waters in the United States, *Radiological Health Data*, 1:20-31 (October 1960).
- (4) Straub, C. P.: Significance of Radioactivity Data, *Journal of the American Water Works Association*, 53:704 (June 1961).
- (5) Setter, L. R., J. E. Regnier, and E. A. Diephaus: Radioactivity of Surface Waters in the United States, *Journal of the American Water Works Association*, 51:1377 (November 1959).
- (6) Robert A. Taft Sanitary Engineering Center, Public Health Service: Radionuclide Analysis of Environmental Samples, Technical Report, R59-6 (1959).
- (7) Straub, C. P., L. R. Setter, A. Goldin, and P. F. Hallbach: Strontium-90 in Surface Waters in the U.S., *Journal of the American Water Works Association*, 52:756 (June 1960).

Radionuclide Analyses of Coast Guard Water Supplies

October 1961–April 1962

U.S. Coast Guard and Division of Radiological Health, Public Health Service

From 1957 through 1960, monthly samples of several U.S. Coast Guard light station rain water cistern supplies were routinely analyzed for beta activity by the Robert A. Taft Sanitary Engineering Center (1, 2). Due to the resultant low radioactivity concentrations two years after the 1958 cessation of nuclear weapons testing, this sampling procedure was terminated.

Since October 1961, drinking water samples have been obtained from Coast Guard Loran Stations in Alaska. In most cases, these samples are collected from small artificial im-

poundments or lakes. The sampling sites are shown in figure 1.

The radionuclide analyses reported in table 1 were performed by the Southwestern Radiological Health Laboratory at Las Vegas, Nevada.

REFERENCES

- (1) Straub, C. P.: Statement on New Data on Uptake in Milk, Food, and Human Bone, *Joint Committee on Atomic Energy Hearings on Fallout From Nuclear Weapons Tests*, 2:990 (May 1959).
- (2) U.S. Coast Guard: Analyses of Cistern Water, *Radiological Health Data*, June 1960, July 1960, and August 1961.

TABLE 1.—RADIONUCLIDES IN WATER SAMPLES FROM U. S. COAST GUARD LORAN STATIONS, OCTOBER 1961–APRIL 1962

[Concentrations in $\mu\text{c}/\text{liter}$]

| Alaskan location | Collection date | Gross alpha | Gross beta | Sr ⁹⁰ | Ra ²²⁶ | I ¹³¹ | Ba ¹⁴⁰ | Cs ¹³⁷ | Zr ⁹⁵ |
|------------------|-----------------|-------------|------------|------------------|-------------------|------------------|-------------------|-------------------|------------------|
| Adak | Dec. 14, 1961 | 5.8 | 12 | 1.5 | ^a ND | ND | ND | ND | ND |
| | Jan. 2, 1962 | 12.2 | 40 | 0.9 | 0.1 | 40 | 40 | ND | 20 |
| | Feb. 13, 1962 | 0.7 | 11 | 0.3 | 0.2 | ND | 20 | ND | ND |
| | Mar. 20, 1962 | ND | 28 | ^b — | — | ND | 15 | ND | ND |
| Attu | Dec. 22, 1961 | ND | 51 | 0.4 | ND | 30 | 30 | 10 | 30 |
| | Jan. 3, 1962 | 3.4 | 85 | 1.2 | 0.8 | ND | 105 | ND | 45 |
| | Feb. 13, 1962 | ND | 80 | 1.7 | ND | ND | 45 | ND | 20 |
| | Mar. 26, 1962 | 0.7 | 40 | 1.7 | 0.1 | ND | 25 | 5 | ND |
| Biorka | Dec. 14, 1961 | 1.5 | 137 | 3.4 | 0.3 | ND | 80 | ND | 50 |
| | Jan. 11, 1962 | 0.6 | 134 | 0.9 | ND | ND | 135 | ND | 85 |
| | Feb. 13, 1962 | ND | 173 | 0.9 | 0.4 | ND | 80 | ND | 70 |
| | Mar. 23, 1962 | 2.1 | 156 | 0.9 | ND | ND | 110 | ND | 120 |
| | Apr. 22, 1962 | ND | 122 | 0.6 | — | ND | ND | ND | 85 |
| Cape Sarichef | Dec. 22, 1961 | 16.8 | 7 | 0.4 | ND | ND | ND | ND | ND |
| | Jan. 3, 1962 | 1.0 | 7 | 0.4 | ND | 45 | 10 | 5 | ND |
| | Feb. 14, 1962 | 1.1 | 13 | ND | — | ND | ND | ND | ND |
| | Mar. 19, 1962 | ND | 61 | — | — | ND | ND | ND | ND |
| | Apr. 20, 1962 | ND | 11 | 0.7 | ND | ND | ND | ND | ND |
| Port Clarence | Oct. 1961 | 1.1 | 48 | 0.8 | ND | ND | ND | ND | ND |
| | Nov. 1961 | ND | 12 | 1.1 | 0.3 | ND | ND | ND | ND |
| | Mar. 19, 1962 | ND | 18 | — | 0.4 | ND | ND | ND | ND |
| | Apr. 23, 1962 | 1.1 | 13 | 0.3 | — | ND | ND | ND | ND |
| Sitkinak | Dec. 21, 1961 | 15.6 | 12 | 0.5 | 0.4 | ND | ND | ND | ND |
| | Jan. 6, 1962 | 4.4 | 76 | 0.3 | 0.6 | ND | 60 | ND | 30 |
| | Feb. 14, 1962 | ND | 120 | 1.7 | ND | ND | 55 | ND | 65 |
| | Mar. 20, 1962 | ND | 53 | 2.0 | 0.3 | ND | 10 | ND | 30 |
| | Apr. 26, 1962 | ND | 39 | 0.7 | — | ND | ND | ND | 15 |
| Spruce Cape | Dec. 13, 1961 | 3.3 | 25 | 2.5 | 0.2 | ND | ND | ND | ND |
| | Jan. 4, 1962 | 1.1 | 70 | 0.8 | 0.4 | 40 | 45 | ND | ND |
| | Feb. 13, 1962 | 2.4 | 465 | 2.7 | ND | ND | 185 | 30 | 230 |
| | Mar. 20, 1962 | 1.0 | 109 | 2.8 | 0.3 | ND | 30 | ND | 20 |
| | Apr. 13, 1962 | 1.0 | 80 | 0.6 | — | ND | 15 | ND | 40 |
| | Apr. 16, 1962 | ND | 608 | 6.2 | — | ND | ND | ND | 605 |

^a ND indicates radionuclide not detectable.

^b Dash indicates result not reported.



FIGURE 1.—COAST GUARD LORAN STATION WATER SAMPLING LOCATIONS

Radioactivity in Drinking Water

DRINKING WATER ANALYSIS PROGRAM 1961

Division of Environmental Engineering and Food Protection, Public Health Service

The Water Section Activity of the Interstate Carrier Branch, Division of Environmental Engineering and Food Protection, PHS, has gathered extensive data on the radioactivity content of water supplies used on interstate carriers such as trains, airplanes, ships, and other conveyances operating in interstate commerce. This work has several objectives, among which are:

1. to determine radioactivity content of interstate carrier water supplies for comparison with the revised Public Health Service Drinking Water Standards (1, 2);
2. to establish the background level as the

basis with which to compare future results from the supplies sampled; and

3. to obtain data for study in connection with the occurrence of certain chronic diseases.

Beginning in November 1960, the project is a continuing one under which it is planned ultimately to sample the 850 U.S. water supplies used by interstate carriers. In many instances, there are several sources for a municipality and each is sampled. When the present water supply sampling for the second group of cities is completed, another group of about 100 cities will be chosen.

Each sample is a 4-liter composite of three 270-ml aliquots a day obtained during a two-week period. The gross alpha, gross beta, and strontium-90 analyses are performed at the Southeastern, Northeastern, and Southwestern Radiological Health Laboratories, operated by the Division of Radiological Health. Table 1 presents partial data from the second sampling.

Previous coverage in *Radiological Health Data*:

Period

1961

1960 and 1961

Issue

May 1962

August 1962

REFERENCES

- (1) Public Health Service: Drinking Water Standards, 1962 Revision, PHS Regulations, (March 6, 1962).
- (2) Drinking Water Standards for Radioactivity, *Radiological Health Data*, 3:156-7 (May 1962).

TABLE 1.—RADIOACTIVITY IN DRINKING WATERS

[Concentrations in $\mu\text{C}/\text{liter}$]

| Station | | Population served | Date sampled | Strontium-90 Total solids | Alpha Activity | | | Beta activity | | |
|----------------------|--------|-------------------|----------------|------------------------------|----------------|-----------|-------|---------------|-----------|--------|
| City | State | | | | Suspended | Dissolved | Total | Suspended | Dissolved | Total* |
| Mobile | Ala. | 202,779 | 11/16-12/2/61 | | | | | <3.0 | 6.3 | <9.3 |
| Montgomery | | 134,393 | | | | | | | | |
| Court St. Pump Sta. | | | 5/30-6/13/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Day St. Pump Sta. | | | 5/31-6/15/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| TVA Widows Creek | | | 6/7-6/22/61 | 1.1 | | | | <3.0 | 37.0 | 40.0 |
| Tampa | Fla. | 274,970 | 4/25-5/9/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Atlanta | Ga. | 487,455 | | | | | | | | |
| Chattahoochee Plant. | | | 8/15-9/1/61 | | | | | <3.0 | <3.0 | <6.0 |
| Hemphill Plant. | | | 8/15-8/30/61 | | | | | <3.0 | <3.0 | <6.0 |
| Columbus | | 116,779 | 11/14-11/28/61 | | | | | <3.0 | 4.5 | <7.5 |
| Elkhart | Ind. | 40,274 | 6/16-6/30/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Evansville | | 141,543 | 6/15-6/30/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Muncie | | 68,603 | 1/18-2/1/61 | 0.2 | <0.5 | <0.5 | <1.0 | <2.6 | <2.6 | <5.2 |
| Washington | | 10,846 | 6/13-6/21/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Ashland | Ky. | 31,283 | 1/26-2/9/61 | 0.1 | | <0.5 | | | <2.6 | |
| Paducah | | 34,479 | 11/2-11/17/60 | | | | | <1.0 | 6.8 | 7.8 |
| Louisville | | 390,639 | 6/14-6/28/61 | | <0.5 | <0.5 | <1.0 | <3.0 | 3.2 | <6.2 |
| Portland | Maine | 72,566 | 3/20-4/3/61 | | <0.5 | <0.5 | <1.0 | <3.0 | 3.9 | <6.9 |
| Ann Arbor | Mich. | 67,340 | 4/20-5/6/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Greenville | Miss. | 41,502 | 1/12-1/27/61 | 0.5 | <0.5 | <0.5 | <1.0 | <2.6 | <2.6 | <5.2 |
| Jackson | | 144,422 | 6/1-6/15/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Vicksburg | | 29,130 | 2/15-3/2/61 | 0.6 | | <0.5 | | | 4.3 | |
| Charlotte | N. C. | 201,564 | | | | | | | | |
| Hoskins Filter Plant | | | 3/30-4/14/61 | 1.0 | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Vest Station | | | 3/30-4/14/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Raleigh | | 93,931 | 4/10-4/24/61 | 1.0 | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Winston-Salem | | 111,135 | 2/14-3/1/61 | 0.2 | | <0.5 | | | <2.6 | |
| Akron | Ohio | 290,351 | 8/18-9/1/61 | | | | | 3.6 | <3.0 | <6.6 |
| Lancaster | Pa. | 61,055 | | | | | | | | |
| Conestoga Plant | | | 7/13-7/27/61 | | | | | <3.0 | <3.3 | <6.3 |
| Susquehanna Plant | | | 8/29-9/12/61 | | | | | <3.0 | <3.0 | <6.0 |
| Charleston | S. C. | 65,925 | 1/10-1/25/61 | | | | | <3.0 | <3.0 | <6.0 |
| North Augusta | | 10,348 | 9/4-9/17/61 | | | | | <3.0 | <3.0 | <6.0 |
| Chattanooga | | 130,009 | 7/12-7/28/61 | 3.2 | | | | <3.0 | 45.0 | 48.0 |
| Clarksville | Tenn. | 22,021 | 7/6-7/24/61 | | <0.5 | <0.5 | <1.0 | <3.0 | <3.0 | <6.0 |
| Knoxville | | 111,827 | 8/18-9/1/61 | | | | | <3.0 | <3.0 | <6.0 |
| Huntington | W. Va. | 83,627 | 12/9-12/27/61 | 0.7 | | | | <1.0 | <2.2 | <3.2 |

* Where "less than" values comprise not more than 10 percent of a total, the "less than" sign is not shown with the total.

SECTION V.—OTHER DATA

Whole Body Counting

Whole body counters are being applied where the detection of low levels of radioactivity in the human body is important, as, for example in the evaluation of hazards to radiation workers and the general population. These instruments are also used in human physiological and pathological investigations as well as in studies related to counter design for special purposes. Discussion and presentation of whole body counting results in previous issues of *RHD* have been limited to medical research programs employing thallium-activated sodium iodide crystals or liquid scintillation solutions. As other whole body counting data become available, they will be published.

CESIUM-137 IN MAN

March 1962 through May 1962

U.S. Army Medical Research Unit, Landstuhl, Germany

In 1955, cesium-137 was first detected in man by the Argonne National Laboratory. It emits a 0.661 Mev gamma photon, which can be quantitatively determined by a properly calibrated whole body counter. Since cesium is physiologically similar to potassium, and for

the most part exists intracellularly, the cesium-137 levels are usually expressed in micromicrocuries per gram of potassium. The whole body counting facility at the Medical Research Unit, Landstuhl, Germany, in its program for measuring the cesium-137 levels in man, utilizes a liquid scintillation counter (1).

TABLE 1.—ASSAYS PERFORMED AT U.S. ARMY MEDICAL RESEARCH UNIT, LANDSTUHL, GERMANY

| Date | Number of subjects | Residence | Cesium-137 $\mu\mu\text{C/gK}$ (average) | Percent MPBB* |
|------------|--------------------|--------------|------------------------------------------|---------------|
| March 1962 | 250 | West Germany | 27 | 0.13 |
| April 1962 | 168 | West Germany | 28 | 0.13 |
| May 1962 | 418 | West Germany | 34 | 0.16 |

* Percent maximum permissible body burden was calculated using 3 μC of cesium-137 as the general population maximum permissible whole body burden and 140 grams of potassium in the body of the "standard man." (See pp. 187 and 192 of Radiological Health Handbook, Office of Technical Services, U. S. Department of Commerce (1960), price \$3.75.)

Previous coverage in *Radiological Health Data*:

| Period | Issue |
|----------------------|--------------|
| 1958 and 1959 | October 1960 |
| 1959 and 1960 | January 1961 |
| 1958, 1959, and 1960 | April 1961 |
| First quarter 1961 | July 1961 |
| Second quarter 1961 | October 1961 |
| Third quarter 1961 | January 1962 |
| First quarter 1962 | August 1962 |

REFERENCE

- (1) *Radiological Health Data* 2:192-4 (April 1961).

External Gamma Activity

RADIATION SURVEILLANCE NETWORK

July 1962

Division of Radiological Health, Public Health Service

Daily measurements of external gamma radiation are made at stations of the Radiation Surveillance Network to assure detection of any substantial deviations from normal background levels. Portable Geiger-Mueller survey instru-

ments are used to obtain measurements at three feet above the ground surface. July 1962 data reported in table 1 are characteristic of individual station observations which in recent years have defined the range of background values.

TABLE 1.—EXTERNAL GAMMA ACTIVITY, JULY 1962

| Station location | | Average (mr/hr) | Station location | | Average (mr/hr) |
|------------------|--------|--------------------|------------------|---------|--------------------|
| City | State | | City | State | |
| Adak | Alaska | 0.01 | Minneapolis | Minn. | 0.01 |
| Anchorage | Alaska | 0.01 | Jackson | Miss. | 0.01 |
| Attu | Alaska | 0.01 | Pascagoula | Miss. | 0.01 |
| Fairbanks | Alaska | 0.01 | Jefferson City | Miss. | 0.01 |
| Juneau | Alaska | 0.01 | Helena | Mont. | 0.03 |
| Kodiak | Alaska | 0.01 | Lincoln | Nebr. | 0.08 |
| Nome | Alaska | 0.01 | Las Vegas | Nev. | 0.01 |
| Point Barrow | Alaska | 0.01 | Concord | N. H. | — |
| St. Paul Island | Alaska | 0.01 | Trenton | N. J. | 0.02 |
| Phoenix | Ariz. | 0.01 | Santa Fe | N. Mex. | 0.04 |
| Little Rock | Ark. | 0.01 | Albany | N. Y. | 0.02 |
| Berkeley | Calif. | 0.01 | Buffalo | N. Y. | 0.01 |
| Los Angeles | Calif. | 0.02 | New York | N. Y. | — |
| Denver | Colo. | 0.02 | Gastonia | N. C. | 0.02 |
| Hartford | Conn. | 0.01 | Bismarck | N. D. | 0.01 |
| Dover | Del. | 0.01 | Columbus | Ohio | 0.01 |
| Washington | D. C. | 0.04 | Painesville | Ohio | 0.01 |
| Jacksonville | Fla. | 0.01 | Oklahoma City | Okla. | 0.02 |
| Miami | Fla. | 0.01 | Ponca City | Okla. | 0.02 |
| Atlanta | Ga. | — | Portland | Ore. | 0.02 |
| Agana | Guam | 0.01 | Harrisburg | Pa. | 0.02 |
| Honolulu | Hawaii | 0.01 | San Juan | P. R. | 0.01 |
| Boise | Idaho | 0.01 | Providence | R. I. | 0.02 |
| Springfield | Ill. | 0.01 | Columbia | S. C. | 0.02 |
| Indianapolis | Ind. | 0.01 | Pierre | S. D. | 0.02 |
| Iowa City | Iowa | 0.01 | Nashville | Tenn. | 0.01 |
| Topeka | Kans. | 0.03 | Austin | Tex. | 0.01 |
| Frankfort | Ky. | 0.01 | El Paso | Tex. | 0.02 |
| New Orleans | La. | 0.02 | Salt Lake City | Utah | 0.02 |
| Augusta | Maine | 0.02 | Barre | Vt. | 0.02 |
| Presque Isle | Maine | 0.02 | Richmond | Va. | 0.01 |
| Baltimore | Md. | 0.02 | Seattle | Wash. | 0.02 |
| Rockville | Md. | — | Charleston | W. Va. | — |
| Lawrence | Mass. | 0.02 | Madison | Wis. | 0.01 |
| Winchester | Mass. | 0.01 | Cheyenne | Wyo. | 0.02 |
| Lansing | Mich. | 0.02 | | | |

*Dash indicates no data received.

Environmental Levels of Radioactivity at Atomic Energy Commission Installations

The U.S. Atomic Energy Commission receives from its contractors quarterly reports on the environmental levels of radioactivity in the vicinity of major Commission installations. The reports include data from routine monitoring programs where operations are of such a nature that plant perimeter surveys are required.

Summaries of the environmental radioactivity data for 21 AEC installations have appeared periodically in *Radiological Health Data* since November 1960. Summaries follow for Hanford Atomic Products Operation, Lawrence Radiation Laboratory, and Los Alamos Scientific Laboratory.

The measured concentration of a radionuclide in air and water may be compared with the Maximum Permissible Concentration (MPC) of that nuclide as recommended by the National Committee on Radiation Protection and Measurement (NCRP). For the environment near an AEC installation, the applicable MPC's are one-tenth of the occupational MPC values for continuous exposure given in National Bureau of Standards "Handbook 69." The MPC values applicable to the following reports are given in table 1.

To avoid any possible misunderstandings in

the use of the term "Maximum Permissible Concentrations," the Federal Radiation Council (FRC) has recommended the term "Radioactivity Concentration Guide" (RCG) for the use of Federal Agencies. Although FRC has not published a table of RCG values, several AEC installations, have adopted FRC nomenclature and use the NCRP published values or have derived their own RCG values on the basis of Radiation Protection Guides (RPG's) such as 500 mrem/year total body exposure. For a more detailed description of the meaning and use of the RCG, the reader is referred to FRC Report No. 2.* The RPG's in terms of total intake in $\mu\mu\text{C/day}$ averaged over a 1-year period which are recommended in FRC Report No. 2 are as follows: the upper limits of Range II for radium-226, iodine-131, strontium-90 and strontium-89 are 20, 100, 200, and 2000 respectively.

In the following reports, the use of non-specific terms such as "total activity," "total alpha," and "gross beta" do not in themselves suggest any one MPC value. Often, when con-

* Federal Radiation Council: *Background Material for the Development of Radiation Protection Standards, Report No. 2*, Superintendent of Documents, U.S. Government Printing Office, Washington 25, D.C., price 20 cents.

TABLE 1.—SELECTED ENVIRONMENTAL MPC VALUES PERTAINING TO AEC INSTALLATION REPORTS IN THIS SUBSECTION

| Line No. | Radionuclide or mixture of unknown nuclides | Environmental MPC's | |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|---------------------------------|
| | | Water ($\mu\mu\text{C/liter}$) | Air ($\mu\mu\text{C/m}^3$) |
| 1 | If Sr^{90} , I^{129} , Pb^{210} , Po^{210} , At^{211} , Ra^{223} , Ra^{224} , Ra^{226} , Ac^{227} , Ra^{228} , Th^{230} , Pa^{231} , Th^{232} , and Th-nat are not present ^a | 3,000 | — |
| 2 | If Sr^{90} , Pb^{210} , Ra^{226} , Ra^{228} are not present ^a | 600 | — |
| 3 | If Ra^{226} , Ra^{228} are not present ^a | 100 | — |
| 4 | Mixture of unidentified nuclides | 10 | 0.04 |
| 5 | If α emitters and Ac^{227} are not present ^a | — | 1.0 |
| 6 | If α emitters and Pb^{210} , Ac^{227} , Ra^{223} , Pu^{239} are not present ^a | — | 10 |
| 7 | If α emitters and Sr^{90} , I^{129} , Pb^{210} , Ac^{227} , Ra^{223} , Pa^{231} , Pu^{239} , Bk^{240} are not present ^a | — | 100 |
| 8 | Arsenic-76 | 20,000 | 3,000 |
| 9 | Copper-64 | 200,000 | 40,000 |
| 10 | Chromium-51 | 2,000,000 | 80,000 |
| 11 | Iodine-131 (FRC Guide: 100 $\mu\mu\text{C/day}$) | 2,000 | 300 |
| 12 | Manganese-56 | 100,000 | 20,000 |
| 13 | Neptunium-239 | 100,000 | 30,000 |
| 14 | Phosphorus-32 | 20,000 | 2,000 |
| 15 | Plutonium-239 | 5,000 | 0.06 |
| 16 | Silicon-31 | 200,000 | 30,000 |
| 17 | Sodium-24 | 30,000 | 5,000 |
| 18 | Strontium-90 | 100 | 10 |
| 19 | Uranium-natural | 20,000 | 2 |

^a "Not present" implies that the concentration of the nuclide is small compared with its appropriate MPC. According to recent AEC regulation (Federal Register, Title 10, Part 20, August 9, 1961), a group of nuclides may be considered not present if the ratio of each nuclide to its appropriate MPC is equal to or less than 1/10 and if the sum of these ratios for the group in question is equal to or less than 1/4.

centrations are low a laboratory will assign an MPC value that is more restrictive than necessary. This avoids the more costly isotopic tests necessary to justify a less restrictive value. References to table 1 will be made to designate the appropriate MPC's reported by the laboratories.

HANFORD ATOMIC PRODUCTS OPERATION

Third and Fourth Quarters 1961

*General Electric Company
Richland, Washington*

The Hanford Atomic Products Operation (HAPO) is located in a semi-arid region of southeastern Washington. The project is a five-hundred square mile complex of nuclear reactors, fuel fabrication plants, chemical separation facilities, and research and development laboratories.

Various radioactive wastes which result from plant operations are disposed through controlled releases to the atmosphere, ground, and the waters of the Columbia River. A primary responsibility of plant personnel is to monitor and safeguard against any effects these wastes may have on the 80,000 residents of Richland, Pasco, Kennewick, and other communities in the vicinity of the project.

The monitoring program includes surveillance of the atmosphere, vegetation, fish and wildlife, milk and agricultural produce, and the waters of the Columbia River and Pacific Ocean. Further measurements are made of external radiation and possible ground contamination from burial pits.

Atmosphere

Gaseous wastes are released to the atmosphere through 200-foot stacks after removal of most radioactive material originally present.

Hanford project maintains air sampling stations at Benton City, Kennewick, Pasco, Richland, Seattle, Spokane, Walla Walla, and Yakima, Washington; Meacham and Klamath Falls, Oregon; Boise and Lewiston, Idaho; and Great Falls, Montana. Sample filters are changed weekly and sent to Hanford for analy-

TABLE 2.—AVERAGE AIR-BORNE BETA ACTIVITY IN SELECTED PACIFIC N. W. LOCATIONS, HAPO

[Average concentrations in $\mu\text{C}/\text{m}^3$]

| Sampling location | Third Quarter 1961 | | Fourth Quarter 1961 | |
|----------------------|--------------------|------|---------------------|------|
| | No. of samples | Beta | No. of samples | Beta |
| Richland, Wash.----- | 13 | 0.56 | 14 | 6.10 |
| Boise, Idaho----- | 11 | 0.44 | 13 | 8.70 |
| Hanford project----- | 13 | 0.54 | 14 | 4.70 |
| Seattle, Wash.----- | 13 | 0.18 | 14 | 3.38 |
| Meacham, Ore.----- | 13 | 0.15 | 13 | 4.14 |

ses. Third and fourth quarter 1961 summaries of beta concentration in these filters are presented in table 2.

Releases of iodine-131 are of particular interest and routine measurements are made at several locations around the plant area. Total stack releases averaged 0.78 and 0.55 curies per day for the third and fourth quarters, respectively, and the concentrations in the local environmental air are shown in table 3.

TABLE 3.—IODINE-131 IN AIR AT PERIMETER COMMUNITIES, HAPO

[Average concentration in $\mu\text{C}/\text{m}^3$]

| Station | Third Quarter 1961 | | Fourth Quarter 1961 | |
|----------------|--------------------|------------------|---------------------|------------------|
| | No. of samples | I ¹³¹ | No. of samples | I ¹³¹ |
| Richland | 13 | 0.011 | 13 | 0.054 |
| North Richland | 13 | 0.014 | 13 | 0.118 |
| Benton City | 12 | 0.022 | 12 | 0.026 |
| Pasco | 13 | 0.015 | 11 | 0.107 |

Water

Hanford's eight production reactors are cooled by treated water pumped from the Columbia River. After the water passes through the reactors, it is returned to the river. Some radioactivity results from neutron activation of impurities remaining in the treated water and is discharged in the effluent. The major part (90 percent) of this radioactive material consists of Mn⁵⁶, Cu⁶⁴, Na²⁴, Cr⁵¹, Np²³⁹, As⁷⁶, and Si³¹. Although the presence of Zn⁶⁵, P³², and certain other radionuclides are of interest, they are found only in minor quantities. River water samples are taken at Hanford Ferry, Pasco, and Vancouver to be analyzed for several radionuclides. Selected results are presented in table 4.

TABLE 4.—SELECTED RADIONUCLIDES IN COLUMBIA RIVER WATER, HAPO

[Average concentration in $\mu\mu\text{c/liter}$]

| Sampling location | Third Quarter 1961 | | | Fourth Quarter 1961 | | |
|-------------------|--------------------|-----------------|------------------|---------------------|-----------------|------------------|
| | Na ²⁴ | P ³² | Zn ⁶⁵ | Na ²⁴ | P ³² | Zn ⁶⁵ |
| Hanford Ferry | 9150 | 184 | 350 | 11,600 | 328 | 500 |
| Pasco | 1750 | 99 | 158 | 2,120 | 234 | 238 |
| Vancouver | — | 22 | 87 | — | 42 | <24 |

*Dash indicates no analysis.

Fish and Wildlife

Fish in the Columbia River accumulate some of the radionuclides present in water and whitefish in particular concentrate phosphorus-32 which is a major bone seeker. The average concentration of phosphorus-32 in the flesh of whitefish caught near the plant was 800 $\mu\mu\text{c/g}$.

Migratory waterfowl, which may have utilized the Hanford section of the Columbia River were obtained from hunters in Washington, Oregon, Idaho, and California. Analyses of 1500 of these specimens showed that only one out of thirteen contained concentrations of phosphorus-32 greater than the detection level of 50 $\mu\mu\text{c/g}$ and the maximum concentration found was 2,600 $\mu\mu\text{c P}^{32}/\text{g}$.

In order to determine the kinds, number, and disposition of local fish and waterfowl harvested, a creel survey was initiated in 1961. This survey is being carried out by contract with the Washington Department of Game.

An estimated 10 pounds of whitefish per person per year for roughly 2,000 people in the Pasco-Kennewick-Richland area was used in the evaluation of population dose from fish and waterfowl: (*Editor's note: Refer to Radiological Health Data, February 1962, pp. 63-4 for a graphic presentation of the estimated population dose in the Hanford vicinity for 1960. The 1961 estimates are similar.*)

Milk and Agricultural Produce

There is no farming within about a 20-mile radius of the chemical separations facilities (the major local source of air-borne radionuclides), but the Ringold farms and River-view District of Pasco, which are respectively about 15 and 30 miles downstream from the

reactors, take water from the Columbia River. Some radionuclides are traced though the irrigation process into milk and produce.

Analyses of local milk for strontium-90, strontium-89, and cesium-137 indicated average concentrations which ranked among the lowest in the nation. These radionuclides result predominantly from world-wide fallout. However, phosphorus-32 and zinc-65 in milk originate from the irrigation water and the 1961 average concentrations of phosphorus-32 in milk from dairy farms in the Ringold and Pasco areas were 725 and 622 $\mu\mu\text{c/liter}$ respectively. The zinc-65 average concentration was 725 $\mu\mu\text{c/liter}$ at Ringold and 518 $\mu\mu\text{c/liter}$ at Pasco.

Assays of local milk for iodine-131 during the third quarter revealed little contamination and the average was less than 60 $\mu\mu\text{c/liter}$. Fallout after the resumption of nuclear testing caused a brief rise in iodine-131 and on November 2, 1961 the maximum local concentration reached 1,500 $\mu\mu\text{c/liter}$. However, the rapid decay rates of iodine-131 and the use of dry winter feeds allowed the concentration in milk to drop to about 2 $\mu\mu\text{c/liter}$ by late December.

Measurements were made on samples of pasture grass to estimate the iodine-131 intake from leafy vegetables. Prior to the arrival of world-wide fallout, the average concentration was 0.2 $\mu\mu\text{c/g}$; after the arrival of fallout, external sources of nuclides introduced a large source of error which interfered with estimations of exposure from plant emissions.

External Radiation

Ionization chambers located above ground and in the Columbia River were used to estimate the combined exposure from external sources in the Hanford area. Aboveground measurements indicated an annual exposure in 1961 of about 130 mr. Virtually all of this radiation is attributable to natural background.

The presence of gamma emitters, especially Na²⁴, in reactor effluent causes a higher exposure rate in the river. The average dose in the water near Richland was about 2.5 mr/day and near Kennewick about 0.9 mr/day. This would mean that a person swimming or boating in the river for 240 hours per year would receive about 25 mr of exposure.

Ground Releases

Solid wastes are packaged and buried in trenches; "high level" liquid wastes which normally contain activity of 10^{11} μc /liter or more are neutralized and stored in concrete tanks lined with steel; "intermediate level wastes" containing activity between 5×10^4 and 10^{11} μc /liter are sent to underground cribs from which waste can percolate into the soil; and "low level wastes" containing less than 5×10^4 μc /liter are sent to depressions in the ground where they form surface ponds.

To monitor the extent to which radionuclides reach the ground water, wells have been drilled in crib and tank storage areas. Figure 1 shows the probable extent and concentration of radioactive materials in ground water.

Previous coverage in *Radiological Health Data*:

| Period | Issue |
|-------------------------------------------------------------------|---------------|
| 1959 and first and second quarters 1960 | May 1961 |
| Third and fourth quarters 1960 and first and second quarters 1961 | February 1962 |

LAWRENCE RADIATION LABORATORY Third and Fourth Quarters 1961

University of California
Berkeley, California
Berkeley Site

The Berkeley site of the Lawrence Radiation Laboratory (LRL) is located adjacent to the campus of the University of California. The technical facilities include a 6.3 Bev proton accelerator, a 700 Mev cyclotron, a linear accelerator of 10 Mev per nucleon, and various chemistry and physics laboratories.

The environmental sampling program includes atmospheric samples of stack releases, local area air, perimeter air, and rain or dry deposition; and water samples of building wastes, tap water, surface water, and off-site creeks. Analyses are made for alpha and beta concentrations.

Approximately 30 to 40 stacks, mostly laboratory hood exhausts, were sampled during the 3rd and 4th quarters. The normal sampling period was one week. Filters were then counted for alpha and beta activity.

Outdoor air samples were taken daily or weekly at three locations within the laboratory area and at four locations along the University of California property line. One additional sampling site is now located in downtown Berkeley to provide background information on contamination from weapons testing.

A summary of 3rd and 4th quarter air monitoring data is presented in table 5.

TABLE 5.—ATMOSPHERIC MONITORING,
BERKELEY SITE

(Average concentration in $\mu\text{c}/\text{m}^3$)

| Sampling locations | Third Quarter 1961 | | | Fourth Quarter 1961 | | |
|-------------------------------|--------------------|--------|------|---------------------|--------|------|
| | No. of samples | Alpha | Beta | No. of samples | Alpha | Beta |
| Stacks | 498 | 0.017 | 0.60 | 404 | 0.031 | 5.30 |
| Local air | 70 | 0.002 | 0.82 | 85 | <0.002 | 8.43 |
| Perimeter air | *16 | <0.002 | 2.40 | 47 | <0.002 | 8.29 |
| Reference sample ^b | — | — | — | 32 | 0.003 | 6.35 |

^a Predominantly September samples.

^b Downtown Berkeley location began operation November 1961.

Rain and/or dry deposition was sampled at irregular intervals at one location during the 3rd quarter, but beginning with the 4th quarter the sampling program was increased to provide weekly data at 7 sites. Fourth quarter deposition averages may be found in table 6.

TABLE 6.—TOTAL DEPOSITION AT BERKELEY SITE
FOURTH QUARTER 1961

(Deposition in $\text{m}\mu\text{c}/\text{m}^2$)

| Sampling location | No. of samples | Alpha | Beta |
|-------------------|----------------|-------|------|
| Local area | | | |
| Building 4 | 16 | 0.170 | 467 |
| All others | 24 | 0.024 | 153 |
| Perimeter | 47 | 0.034 | 168 |

Acid wastes from two chemical laboratory buildings were sampled and analyzed before dilution and discharge through the main sewer system. The sewage data in table 7 were estimated from the results of the above analyses. It is assumed that these two buildings are the only significant sources of radioactive liquid wastes.

Surface water was sampled weekly or bi-weekly from two small streams on the laboratory site. Six nearby off-site creeks were also sampled every two or four weeks depending on the relative distance from the laboratory.

Tap water samples were taken approximately every week and the analyses results are summarized in table 7 with the above water data.

TABLE 7.—WATER MONITORING, BERKELEY SITE

[Average concentration in $\mu\text{c}/\text{liter}$]

| Type of sample | Third Quarter 1961 | | | Fourth Quarter 1961 | | |
|--------------------|--------------------|-------|------|---------------------|-------|------|
| | No. of samples | Alpha | Beta | No. of samples | Alpha | Beta |
| Tap water | 10 | 0.04 | 2.0 | 11 | 0.04 | 5.1 |
| On site streams | 24 | 0.12 | 7.9 | 24 | 0.14 | 21.1 |
| Off site streams | 33 | 0.10 | 2.6 | 30 | 0.09 | 12.5 |
| Sewage (estimated) | — | 0.09 | 3.6 | — | 0.15 | 10.3 |

Livermore Site

The Livermore site of LRL is located 3 miles east of Livermore, California. Technical facilities include a small cyclotron, a two megawatt swimming pool reactor, and physics and chemistry operations associated with a weapons development program. The laboratory also operates Site 300 which is physically located about 20 miles southeast of Livermore.

Environmental survey samples are collected for the purpose of detecting changes from the normal concentrations of radionuclides that could be attributed to operations at the laboratory. Samples include air particulate, soil, water, and sewage plant products.

Air particulate samples are collected continuously from 11 sites within 5 miles of the LRL. Filters are changed weekly and analyses made for alpha and beta activity. Third and fourth quarter results of these locations are summarized in table 8. The third quarter averages of alpha and beta activity at Site 300 were <0.0014 and $<0.0032 \mu\text{c}/\text{m}^3$, respectively.

TABLE 8.—AIR MONITORING, LIVERMORE SITE

[Average concentration in $\mu\text{c}/\text{m}^3$]

| Distance from lab. | Third Quarter 1961 | | | Fourth Quarter 1961 | | |
|--------------------|--------------------|-------|-------|---------------------|-------|------|
| | No. of samples | Alpha | Beta | No. of samples | Alpha | Beta |
| 0.5-1.0 miles | 13 | 0.012 | 0.039 | 44 | 0.006 | 3.4 |
| 2-5 miles | 13 | 0.012 | 0.045 | 135 | 0.007 | 3.2 |

Ground water samples are collected to insure that radioactivity is not entering underground water sources from laboratory operations. This

underground water is the sole source of domestic water for the Livermore area and is also used for irrigation. All water used at Site 300 is drawn from an underground source which is geologically separated from Livermore water. Table 9 summarizes the quarterly results.

TABLE 9.—RADIOACTIVITY IN WELL WATER AT LIVERMORE SITE

[Average concentration in $\mu\text{c}/\text{liter}$]

| Sample source | Third Quarter 1961 | | | Fourth Quarter 1961 | | |
|------------------|--------------------|-------|--------|---------------------|-------|------|
| | No. of samples | Alpha | Beta | No. of samples | Alpha | Beta |
| LRL Site | 65 | <15 | <100 | 51 | 8.2 | 68 |
| Site 300 | 65 | <15 | <100 | 30 | 8.2 | 68 |
| Livermore (city) | 6 | <15 | <100 | 8 | 8.2 | 68 |

Numerous soil samples are taken from around the Livermore area to maintain a further check on air-borne radioactive particulates. Sewage from the laboratory and Livermore is processed into dry sludge used for agricultural soil conditioner. Therefore, samples are taken at the Livermore Sewage Treatment Plant to assure that the concentration of radioactivity in the sludge is no higher than the normal concentration in the soil. Results of soil and sewage sludge sampling are presented in table 10.

TABLE 10.—RADIOACTIVITY IN SOIL AND SLUDGE, LIVERMORE SITE

[Average concentration in $\mu\text{c}/\text{g}$]

| Sample | Third Quarter 1961 | | | Fourth Quarter 1961 | | |
|---------|--------------------|--------|-------|---------------------|-------|------|
| | No. of samples | Alpha | Beta | No. of samples | Alpha | Beta |
| Soil | 39 | <7.7 | <52 | 54 | 4.5 | 38 |
| Sludge* | 10 | <7.7 | <52 | 3 | 4.5 | 38 |

*Livermore Sewage Plant.

LOS ALAMOS SCIENTIFIC LABORATORY 1960-1961

University of California
Los Alamos, New Mexico

The Los Alamos Scientific Laboratory (LASL), in cooperation with the U.S. Geological Survey, conducts a program of environmental monitoring to determine what effect

Laboratory activities have on the local area. The Laboratory is located in an area of high elevation in northern New Mexico. The program includes air particulate sampling for alpha and beta activity, water assays for specific radioactive materials, and measurements of gamma radiation. Average annual rainfall is 18 inches and stream flow is seasonal, dependent on snow run-off and storm activity.

Radioactive wastes are released in small amounts to the atmosphere and stream beds or are buried in restricted burial areas. Disposal is governed by the geology of the area to minimize any possibility of long-lived radioactive material causing future contamination of drinking or irrigation water sources.

Air Monitoring

Daily sampling of beta activity in air at LASL showed results similar to the values reported throughout the United States. The 1961 average at the laboratory was $3.32 \mu\mu\text{C}/\text{m}^3$. This may be compared with $4.22 \mu\mu\text{C}/\text{m}^3$ for the 1961 average of 171 stations of the Public Health Service National Air Sampling Network.¹

Air particulate sampling stations are also located at 24 sites throughout the area to monitor for long-lived alpha activity. Sampling is performed continuously for one week before the filters are removed and counted. The maximum reading at any station for one week during the two-year period was $0.02 \mu\mu\text{C}/\text{m}^3$; the average was less than $0.004 \mu\mu\text{C}/\text{m}^3$ (See table 1, line 4 for MPC).

Water Monitoring

Assays for plutonium and uranium are made on water samples taken from the Los Alamos environs including surface and ground water, supply wells, and test wells.

The Rio Grande and Rio Chama are sampled periodically. The results indicate a fairly constant uranium concentration of about $1 \mu\mu\text{C}/\text{liter}$. Bi-monthly or semi-annual samples collected from seven intermittent surface water sources showed uranium concentrations similar to that of the river samples.

Thirteen supply wells and 8 test wells are also sampled. The majority of these wells contain

the same uranium concentration as the river and surface sources. One exception is a deep supply well which has a concentration of about $10 \mu\mu\text{C}/\text{liter}$.

No plutonium has been detected in any water sample taken from the above sources.

Gamma Radiation

Film badges are placed on stakes at approximately 100 locations throughout Los Alamos County to detect and measure any gamma radiation resulting from LASL activities. The badges are presently collected and read once each quarter.

During 1960, two stations received slightly more than 0.5 rem of gamma radiation and in 1961 readings of 7.13, 3.15, 2.00, 0.75, and 0.54 rem were obtained at five stations. However, all of these stations were immediately adjacent to a radiation facility and were located so that personnel could not have remained long enough in the area to receive an appreciable part of the total dose. The source of the above high readings was removed on December 13, 1961 and the measurements indicate that no individual of the population could have received over 0.5 rem of whole body radiation during a 12 month period.

NEVADA TEST SITE DETONATION SURVEILLANCE

Atomic Energy Commission

April 14 Event

A low yield nuclear device was detonated in an underground tunnel chamber at the Nevada Test Site on April 14, 1962. Immediately upon detonation, radioactive steam was vented from the mouth of the tunnel and from a fissure which occurred on the top of the mesa. Aircraft monitoring of the cloud indicated a heading of 20° from ground zero, and, by twenty-two minutes after detonation, the cloud had reached an altitude of 10,000 feet Mean Sea Level (MSL). The cloud held to this bearing, and it was tracked on the ground and in the air.

The U.S. Public Health Service personnel, under contract to the Atomic Energy Commission, conducted extensive off-site monitoring

¹ *Radiological Health Data* 3:113 (April 1962).

TABLE 11.—AIR MONITORING FOLLOWING THE APRIL 14, 1962 EVENT, NTS

| Location | Number of people | Sampling period | | | Radio-isotopes | Average concentration ($\mu\text{mc}/\text{m}^3$) |
|--------------|------------------|------------------------|----------|------|----------------|--------------------------------------------------------|
| | | Duration (hr.-min.) | Ending | | | |
| | | | Date | Time | | |
| Diablo----- | 5 | 6-50 | April 14 | 1650 | Iodine-131 | 525 |
| | | | | | Iodine-135 | 2,000 |
| | | 14-0 | April 15 | 0700 | Iodine-131 | 17.3 |
| | | | | | Iodine-133 | 87 |
| | | | | | Iodine-135 | * ND |
| Currant----- | ^b 50 | 24-0 | April 15 | 0800 | Iodine-131 | 400 |
| | | | | | Iodine-133 | 2,100 |
| | | | | | Iodine-135 | ND |

* ND means not detectable.

^b Migrant farm population varies from 50-200 during summer and fall.

and have determined the following exposures to the off-site population:

At Diablo Maintenance Station (forty-four miles from the test site), five people were present when the cloud passed. They were wearing badges supplied on March 8, 1962. These badges were collected on April 25, 1962, and showed a total exposure of 55 to 65 mr. After subtracting the normal background dose from the total recorded, lifetime doses of approximately 28 to 38 mr were estimated as resulting from this event.

As the cloud was carried northward by prevailing winds, it arrived over Currant (population 50; 115 miles from NTS) at about H + 7½ hours. For a period of 30 minutes, the reading was above background, and the peak reading was 0.5 mr per hour. Air samples indicated the presence of slight but detectable quantities of fresh fission products, such as iodine-131, iodine-133, tellurium-132 and ruthenium-105.

Table 11 illustrates the amount of activity of radioiodine in the air monitored at populated areas. The periods of collection encompass the passage of the gaseous cloud.

May 19 Event

Following an underground nuclear detonation at the Nevada Test Site on May 19, 1962, radioactive gases were released. Gases escaped for a period of twenty minutes. These gases formed into a cloud and were carried north of the test site by the prevailing winds.

From off-site data collected, the highest measured gamma radiation in a location where people were known to be present was at Diablo, fifty-five miles north of the test site. The cloud arrived at H + 1¾ hours, with a maximum ground reading of 1.5 mr/hr at H + 2½. By H + 3½, the reading was 0.5 mr/hr, and at H + 7¾, readings were at background levels. The total exposure might have been about 1 mr, but this was considerably below the threshold of detection by the film badges worn.

Due to high surface wind speeds, this cloud was carried rapidly northward and had dispersed considerably by the time it passed to the west of Ely. Between 2:00 p.m. and 3:45 p.m., it was scarcely detectable above normal background.

Analysis of activated charcoal filters indicated the atmospheric concentrations of radioisotopes at the time of cloud passage shown in table 12.

Addenda

Additional information now is available from data collected after the March 5, 1962 event, a subsurface nuclear detonation at the Nevada Test Site that released activity to the atmosphere*. These data show that the highest concentrations found in air in a populated place were at Warm Springs, Nevada. The air sampler operating between 1020 and 1515 hours indicated a gross beta count of 1000 $\mu\text{mc}/\text{m}^3$,

* These data are supplemental to the material appearing in *RHD*, May 1962, page 170.

TABLE 12.—AIR MONITORING FOLLOWING THE MAY 19, 1962 EVENT, NTS

| Location | Sampling period | | | Radio-isotopes | Average concentration ($\mu\text{mc}/\text{m}^3$) |
|---------------|------------------------|--------|------|---------------------------------------------------------|--------------------------------------------------------|
| | Duration (hr.-Min.) | Ending | | | |
| | | Date | Time | | |
| Tempiute..... | 7-00 | May 19 | 1630 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ^a ND ^b — ND ND |
| Penoyer..... | 7-35 | May 19 | 1635 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ND 3.7 6.2 ND |
| Diablo..... | 6-15 | May 19 | 1540 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ND 3.2 12 ND |
| Diablo..... | 18-45 | May 20 | 1030 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ND 0.69 1.46 ND |
| Fallini..... | 24-00 | May 20 | 0615 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ND ND ND ND |
| Currant..... | 10-45 | May 20 | 0645 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | 516 23 18 ND |
| Lund..... | 5-30 | May 19 | 1630 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ND ^b — ND ND |
| Ely..... | 5-00 | May 19 | 1400 | Iodine-131 Iodine-133 Iodine-135 Tellurium-132 | ND ND ND ND |

• ND means not detectable.

b Detected but not quantitated.

16 $\mu\mu\text{c}/\text{m}^3$ of tellurium-132, 6.9 $\mu\mu\text{c}/\text{m}^3$ of iodine-131, 187 $\mu\mu\text{c}/\text{m}^3$ of iodine-133 and 275 $\mu\mu\text{c}/\text{m}^3$ of iodine-135 on the filter and 33.7 $\mu\mu\text{c}/\text{m}^3$ of iodine-133 and 73.5 $\mu\mu\text{c}/\text{m}^3$ of iodine-135 on the charcoal cartridge.

Based on dose rate readings, the estimated maximum exposure received off-site was ap-

proximately 1.7 mr to two persons at Clark's Station. (Film badges have a threshold of recording at 30 milliroentgens, and it is difficult to estimate accurately such low doses from survey meter readings.) The internal exposure due to inhalation was a small fraction of the total external exposure.

Reported Nuclear Detonations

September 1962

Summary information on announced nuclear detonations during the last month is indicated in the table below. Not all detected U.S.S.R. tests are necessarily announced by the Atomic Energy Commission. The AEC has indicated in general announcements that additional detonations have been detected in both the Fall 1961 U.S.S.R. test series and during the recent U.S.S.R. testing. The AEC announced by specific press releases 31 tests in the Fall 1961 series, and indicated in more recent press releases additional tests which brought the series' total to approximately 50. During the current U.S.S.R. testing activities, a similar reporting procedure is being used by AEC.

Radiological Health Data summarizes all available specific announcements of nuclear

tests during each month preceding publication. Numbers are assigned to only those U.S.S.R. tests which have been announced since the termination of the test moratorium in 1961.

Corrections for the July 1962 Reported Nuclear Detonations: (1) U.S. test number 67 (Project Sedan) on July 6 at the Nevada Test Site was incorrectly reported as having a yield range of 1,000 kilotons; the correct yield range for this test is 100 kilotons. "Project Sedan" is part of the Commission's Plowshare Program in which peaceful uses for nuclear explosives are being explored. (2) The July 9 high altitude U.S. test has been called "Starfish" and its yield range previously given as megaton is specified as 1.4 megatons.

| Test number | Location | Date (1962) | Yield range* | Type of test |
|-------------|----------|-------------|--------------|--------------|
|-------------|----------|-------------|--------------|--------------|

REPORTED U.S.S.R. DETONATIONS

| | | | | |
|---------|---------------|-------------|-----------------------|-------------|
| 41..... | Novaya Zemlya | September 2 | Intermediate | Atmospheric |
| 42..... | Novaya Zemlya | 8 | Megaton | Atmospheric |
| 43..... | Novaya Zemlya | 15 | Low megaton | Atmospheric |
| 44..... | Novaya Zemlya | 16 | Low megaton | Atmospheric |
| 45..... | Novaya Zemlya | 18 | Low megaton | Atmospheric |
| 46..... | Novaya Zemlya | 19 | Multi-megaton | Atmospheric |
| 47..... | Novaya Zemlya | 21 | Low megaton | Atmospheric |
| 48..... | Novaya Zemlya | 25 | 30 megatons | Atmospheric |
| 49..... | Novaya Zemlya | 27 | Less than 30 megatons | Atmospheric |

REPORTED U.S. DETONATIONS

| | | | | |
|---------|------------------|--------------|-----------|-------------|
| 79..... | Nevada Test Site | September 14 | Low yield | Underground |
| 80..... | Nevada Test Site | 20 | Low yield | Underground |
| 81..... | Nevada Test Site | 29 | Low yield | Underground |

* Low yield range has been announced as less than 20 kilotons yield; intermediate yield is in the range of 20 kilotons up to 1 megaton; and low megaton yield is in the range from 1 to several megatons.

Announcement of Current Training Activities

Public Health Service

November 1962-May 1963

Short courses, designed primarily for public health personnel interested in acquiring basic information in radiological health, are again being offered by the Division of Radiological Health, Public Health Service. These courses cover a broad range of subjects dealing with the prevention and control of radiation hazards.

Extending from one week to ten-weeks duration, courses are given at the Robert A. Taft Sanitary Engineering Center (SEC), Cincinnati, Ohio; the Radiological Health Laboratory

(RHL), Rockville, Maryland; the Southwestern Radiological Health Laboratory (SWRHL), at Las Vegas, Nevada; the Southeastern Radiological Health Laboratory (SERHL), Montgomery, Alabama; and the Northeastern Radiological Health Laboratory (NERHL), Winchester, Massachusetts.

Below is a list of courses available for the remainder of the training year (course numbers in parentheses):

| | |
|----------------------|----------------------------------------------------------|
| Nov. 5-16, 1962 | Occupational Radiation Protection (212) SEC |
| Nov. 26-30, 1962 | Radioactive Pollutants in Water (195) SEC |
| Nov. 26-Dec. 7, 1962 | Basic Radiological Health (211) RHL |
| Dec. 3-7, 1962 | Radionuclides in Water (222) SEC |
| Dec. 3-14, 1962 | Basic Radiological Health (211) SERHL |
| Dec. 10-14, 1962 | Measurement of Airborne Radioactivity (417) SEC |
| Dec. 10-14, 1962 | Medical X-ray Protection (213) RHL |
| Jan. 7-Mar. 15, 1963 | Engineering Aspects of Radiological Health (231) SEC |
| Jan. 7-18, 1963 | Basic Radiological Health (211) SEC |
| Jan. 7-18, 1963 | Medical Aspects of Radiological Health (201) RHL |
| Jan. 21-Feb. 1, 1963 | Occupational Radiation Protection (212) SEC |
| Jan. 28-Feb. 8, 1963 | Basic Radiological Health (211) RHL |
| Feb. 4-22, 1963 | Environmental Radiation Surveillance (244) SEC |
| Feb. 25-Mar. 7, 1963 | Reactor Safety and Hazards Evaluation (233) SEC |
| Mar. 11-15, 1963 | Medical X-ray Protection (213) SEC or RHL |
| Feb. 11-15, 1963 | Medical X-ray Protection (213) RHL |
| Feb. 11-22, 1963 | Basic Radiological Health (211) SWRHL |
| Feb. 25-Mar. 1, 1963 | Management of Nuclear Emergencies (234) SWRHL |
| Mar. 18-May 24, 1963 | Engineering Aspects of Radiological Health (231) SEC |
| Mar. 18-29, 1963 | Basic Radiological Health (211) SEC |
| April 1-12, 1963 | Occupational Radiation Protection (212) SEC |
| April 15-May 3, 1963 | Environmental Radiation Surveillance (244) SEC |
| May 6-16, 1963 | Reactor Safety and Hazards Evaluation (223) SEC |
| May 6-16, 1963 | Basic Radiological Health (211) SEC |
| May 20-24, 1963 | Medical X-ray Protection (213) SEC or RHL |
| Mar. 18-29, 1963 | Basic Radiological Health (211) RHL |
| April 1-5, 1963 | Medical X-ray Protection (213) RHL |
| April 15-26, 1963 | Basic Radiological Health (211) RHL |
| April 29-May 3, 1963 | Medical X-ray Protection (213) RHL |
| May 13-17, 1963 | Radiological Health for Nurses (203) RHL |
| May 20-24, 1963 | Radionuclides in Foods (340) SEC |
| To be announced | Sanitary Engineering Aspects of Nuclear Energy (225) SEC |

For further information on SEC courses, contact the Chief, Training Program, Robert A. Taft Sanitary Engineering Center. Additional details on courses offered at Rockville, Maryland; Las Vegas, Nevada; Winchester, Massachusetts; and Montgomery, Alabama, are available from the Chief, Training Branch, Division of Radiological Health, 1901 Chapman Avenue, Rockville, Maryland.

ern
at
io-
nt-
io-
in-
the
m-

For the

cpm...
dpm...
g...
kg...
km²...
kvp...
m²...
ma...
ml...
Bev...
Mev...
mi²...
ml...
mm...

mm...
mrad...
mrem...
mr/hr...
nc...
nc/m²...

pc...
 μ sc...

O

PI

CI

PI

A.
ad;
om
le,

ata

Western
(L), at
Radio-
Mont-
Radio-
Win-

or the
num-

rt A.
land;
from
ville,

Data

UNITS AND EQUIVALENTS

For the convenience of the *Radiological Health Data (RHD)* reader a selected list of units and equivalents is presented below.

| Symbol | Name | Equivalent |
|-------------------|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| cpm | count per minute | |
| dpm | disintegration per minute | |
| g | gram | |
| kg | kilogram | 1 kg = 1000 gm = 2.2 pounds |
| km ² | square kilometer | |
| kvp | kilovolt peak | |
| m ³ | cubic meter | 1 m ³ = 1000 liters |
| ma | milliampere | |
| ml | milliliter | |
| Bev | billion electron volts | |
| Mev | million electron volts | |
| mi ² | square mile | |
| ml | milliliter | |
| mm | millimeter | precipitation: $\text{mm} = \frac{\mu\text{mc}/\text{m}^2}{\mu\text{mc}/\text{liter}} = \frac{\text{liter}}{\text{m}^2}$ thickness: 1/1000 meter |
| mm | millimeter | |
| mrad | millirad | |
| mrem | millirem | |
| mr/hr | milliroentgen per hour | |
| nc | nanocurie | 1 nc = 1000 pc = 1 mμc = 10 ⁻⁹ curies |
| nc/m ² | nanocurie per square meter | 1 nc/m ² = 1 mμc/m ² = 1,000 μmc/m ² = 1 mc/km ² = 2.59 mc/mi ² |
| pc | picrocurie | 1 pc = 1 μmc = 10 ⁻¹² curies |
| μmc | micromicrocurie | 1 μmc = 2.22 dpm |

ORDER BLANK FOR RHD

To: Superintendent of Documents
Government Printing Office
Washington 25, D.C.

Please enter my Subscription for Radiological Health Data. I am enclosing Money Order ☐
Check ☐ for this subscription. (\$5.00 a year; \$1.50 additional for foreign mailing.)

Please address the RHD as follows: _____



UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON 25, D.C. 1961

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington 25, D.C.

Subscription price \$5.00 a year, \$1.50 additional for foreign mailing.

Price for a single copy of this issue is 50 cents.

U. S. DEPARTMENT OF
HEALTH, EDUCATION AND WELFARE
PUBLIC HEALTH SERVICE
WASHINGTON 25, D. C.

OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U. S. DEPARTMENT OF H. E. W.

If you do not desire to continue receiving this publication, please CHECK HERE ☐;
tear off this label and return it to the above address. Your name will then be
promptly removed from the appropriate mailing list.